arXiv:1212.1481v1 [math.GT] 6 Dec 2012

Word length statistics for Teichmüller geodesics and singularity of harmonic measure

Vaibhav Gadre Joseph Maher Giulio Tiozzo

December 10, 2012

Abstract

Given a measure on the Thurston boundary and a basepoint in Teichmüller space, one can pick a geodesic ray joining the basepoint to a randomly chosen point on the boundary. In particular, we consider two families of measures: the ones which belong to the Lebesgue or visual measure class, and harmonic measures for random walks on the mapping class group generated by a finitely supported distribution. For any geodesic ray, we compute the ratio between the word metric and the relative metric of approximating mapping class group elements, and prove that this ratio tends to infinity along Lebesgue-typical geodesic, while the limit is finite along geodesics which are typical with respect to harmonic measure. As a corollary, we establish singularity of harmonic measure. The same proof works for Fuchsian groups with a cusp.

Contents

1 Introduction **2**
1.1 The case of $PSL(2,\mathbb{Z})$ 4
1.2 Outline 7
1.3 Acknowledgements 9

2 Preliminaries **9**
2.1 Notation 9
2.2 Quadratic differentials and Teichmüller discs 9
2.3 Curve complex and subsurface projections 10
2.4 Short curves and short markings 12
2.5 Projections and isolated intervals 13
2.6 Excursions and twist parameter 14
2.7 Coarse monotonicity for the word metric 18
2.8 Projection to closest Teichmüller lattice point 19

3 Lebesgue measure sampling **20**
3.1 Metric cylinders with bounded area 20
3.2 Asymptotic length of short curves 21
3.3 Average excursion 24
3.4 The word metric 25

4 Hitting measure sampling **26**
4.1 Distance between geodesic and sample path 28
4.2 Intermediate times 31

5 Fuchsian groups **32**

1 Introduction

Let G be a finitely generated group acting properly discontinuously on a metric space X, with basepoint x_0 and boundary ∂X. In many cases, the points on the boundary ∂X can be identified with geodesic rays starting at x_0. For instance, if X is a Riemannian manifold, then the geodesic rays based at x_0 may be identified with the unit tangent space at x_0, and the rotationally invariant measure on the tangent space at x_0 is often called *visual measure*. It then makes sense to consider the behaviour of a "typical" geodesic in the space, chosen according to this measure. We may also consider random walks on X, by taking a random walk on G, and looking at its image in X under the orbit map $g \mapsto gx_0$. If random walks converge to ∂X almost surely, then this gives an alternative measure on ∂X, known as *hitting measure* or *harmonic measure*. A harmonic measure can be pulled back to the unit tangent ball at x_0 using its identification with ∂X. Again, one may consider the behaviour of a "typical" geodesic ray from x_0 with respect to harmonic measure.

Here, we consider some cases in which the quotient $G\backslash X$ has finite volume but is not compact; namely, nonuniform lattices in $SL(2,\mathbb{R})$ and mapping class groups of surfaces of finite type. In these cases, we show that a geodesic typical with respect to one measure exhibits geometrically different behaviour from a geodesic typical with respect to the other measure. The underlying geometric idea is that the quotient $G\backslash X$ contains a cusp, and Lebesgue-typical geodesics penetrate more deeply into the cusp than typical ones with respect to harmonic measure. In order to make the statement precise, we will reason in terms of the ratio between *word metric* and *relative metric*.

Let γ be a geodesic ray from the basepoint x_0, and γ_t a point at distance t from the basepoint along γ. For each time t, let h_t be a group element such that $h_t x_0$ is a closest element of the G-orbit of x_0 to γ_t. We may measure distance in G by the word metric d_G in G, or alternatively by a non-proper relative metric d_{rel}, in which any distance in a subgroup fixing a cusp has constant length, and which we shall describe more precisely in the cases we consider. We shall consider the ratio of these two distances for elements h_t along a typical geodesic, i.e.

$$\rho = \lim_{t \to \infty} \frac{d_G(1, h_t)}{d_{rel}(1, h_t)}.$$

As we shall see, this limit exists along a full measure set of geodesics in either measure. We shall show that the limit is finite for almost all geodesics in harmonic measure, and infinite for almost all geodesics in Lebesgue measure.

We now state precise results for the two cases; then, we discuss the group $PSL(2,\mathbb{Z})$ acting on hyperbolic plane as a concrete, simple example of this phenomenon, and follow with a more detailed outline of the arguments in the general case.

Let G be a Fuchsian group, i.e. a discrete subgroup of $SL(2,\mathbb{R})$, and let X be the hyperbolic plane $\mathbb{H}^2$. Assume that the quotient orbifold $G\backslash\mathbb{H}^2$ has finite area but is non-compact. Given a basepoint x_0, we may identify the unit tangent bundle with the circle $S^1 = \partial\mathbb{H}^2$ at infinity, and the measure induced on the boundary is Lebesgue measure on the unit circle. By [Fur63], a random walk on G projected to $\mathbb{H}^2$ by the orbit map $g \to gx_0$ converges almost surely to the boundary, defining a harmonic measure ν on S^1.

Let d_G be a word metric on the group G with respect to some finite generating set. We shall choose d_{rel} to be an improper word metric on G with respect to an infinite generating set consisting of the given generating set, together with a choice of parabolic subgroup stabilizing a cusp, for each cusp of the quotient $G \backslash \mathbb{H}^2$.

Theorem 1.1. *Let $G < SL_2(\mathbb{R})$ be a Fuchsian group such that the quotient $G \backslash \mathbb{H}^2$ is a non-compact, finite area hyperbolic orbifold. Given a geodesic ray γ starting at the basepoint x_0, let*

$$\rho(\gamma) = \lim_{t \to \infty} \frac{d_G(1, h_t)}{d_{rel}(1, h_t)},$$

where $h_t x_0$ is a closest element of the G-orbit of x_0 to γ_t. Let Leb *be Lebesgue measure on the circle at infinity, and let ν be the harmonic measure determined by a random walk generated by a finitely supported probability measure on G whose support generates G. Then there is a constant $c > 0$ such that*

$$\rho(\gamma) = \begin{cases} \infty & \textit{for } \mathrm{Leb}\textit{-almost all geodesics } \gamma \\ c & \textit{for } \nu\textit{-almost all geodesics } \gamma. \end{cases}$$

In particular, as a corollary we have:

Corollary 1.2. *The harmonic measure ν is singular with respect to Lebesgue measure on S^1.*

Guivarc'h and LeJan [GLJ90], [GLJ93] proved the singularity result for the special case of the congruence subgroup $\Gamma(2)$ of $PSL(2, \mathbb{Z})$ by studying the asymptotic winding around the cusp of the geodesic flow on $\Gamma(2) \backslash \mathbb{H}^2$. The statistic $\rho(\gamma)$ that we study is similar in spirit to asymptotic winding: the underlying reason for the singularity of ν is the fact that ν-typical geodesics wind around cusps less than Lebesgue-typical geodesics.

The previous observation is the starting point for the main result of the paper, namely the generalization of Theorem 1.1 to the case of G being the mapping class group of a surface S of finite type, which acts on the Teichmüller space $\mathcal{T}(S)$ of hyperbolic metrics on S. The Teichmüller metric on $\mathcal{T}(S)$ is preserved by the action of the mapping class group, and the quotient $\mathcal{M}(S) = G \backslash \mathcal{T}(S)$ has finite volume and is non-compact.

We shall use Thurston's compactification of Teichmüller space, the space of *projective measured foliations* $\mathcal{PMF}$, as a boundary for $\mathcal{T}(S)$. There is a natural *Lebesgue measure class* Leb on $\mathcal{PMF}$ given by pulling back Lebesgue measure from the charts defined using train track coordinates. Kaimanovich and Masur [KM96] showed that if μ is a probability distribution on G, whose support generates a non-elementary subgroup, then the image of a random walk on G under the orbit map $g \mapsto gX_0$ converges to a point in $\mathcal{PMF}$ almost surely. We let ν be the corresponding hitting measure.

The set of uniquely ergodic foliations has full measure with respect to both Leb and ν, and so with respect to either measure we may identify a full measure set of points in $\mathcal{PMF}$ with Teichmüller geodesic rays from a basepoint X_0.

The mapping class group is finitely generated, and we shall write d_G for a choice of word metric on G. We shall let d_{rel} be the word metric with respect to an infinite generating set, consisting of adding to a finite generating set the stabilizers of simple closed curves α_i, where the α_i are a set of representatives for orbits of simple closed curves under G. The relative metric is also quasi-isometric to distance in the projection to the curve complex. Let $\mathcal{T}_\epsilon$ be the ϵ-thin part of Teichmüller space, i.e. the set of surfaces which contain a simple closed curve of hyperbolic length at most ϵ. In this case, we restrict to taking limits over points γ_t which do not lie in the thin part $\mathcal{T}_\epsilon$.

Theorem 1.3. *Let G be the mapping class group of a surface of finite type, and let $\mathcal{T}(S)$ be the Teichmüller space of S. Let $\mathcal{T}_\epsilon$ be the thin part of Teichmüller space, for some $\epsilon > 0$ sufficiently small, and fix a basepoint $X_0 \notin \mathcal{T}_\epsilon$. Given a geodesic ray γ starting at X_0, let*

$$\rho(\gamma) = \lim_{\substack{t\to\infty \\ \gamma_t \notin \mathcal{T}_\epsilon}} \frac{d_G(1, h_t)}{d_{rel}(1, h_t)},$$

where $h_t X_0$ is a closest element of the G-orbit of X_0 to γ_t. Let Leb *be a measure on $\mathcal{PMF}$ in the Lebesgue measure class, and let ν be the harmonic measure determined by a random walk generated by a finitely supported probability measure on G whose support generates a non-elementary subgroup of G. Then there is a constant $c > 0$ such that*

$$\rho(\gamma) = \begin{cases} \infty & \textit{for } \text{Leb}\textit{-almost all geodesics } \gamma \\ c & \textit{for } \nu\textit{-almost all geodesics } \gamma. \end{cases}$$

As a corollary, we give a geometric proof of:

Theorem 1.4 ([Gad09])**.** *The harmonic measure ν is singular with respect to Lebesgue measure.*

The proof in [Gad09] is based on the study of train track splittings on $\mathcal{PMF}$. In this paper, instead, we get the Lebesgue measure statistics by using the ergodicity of the Teichmüller geodesic flow, combined with estimates on the volume of the thin part of the space of quadratic differentials. The statistics for harmonic measure follows from linear progress in the relative metric, combined with sublinear tracking between geodesics and sample paths. In the rest of the introduction, we shall analyse in greater detail the strategy of proof.

1.1 The case of $PSL(2, \mathbb{Z})$

We start by describing a concrete classical example of this phenomenon, in which the space X is the hyperbolic plane $\mathbb{H}^2$, thought of as the universal cover of the finite area orbifold arising from the action of $SL(2, \mathbb{Z})$ on $\mathbb{H}^2$.

The group $SL(2, \mathbb{Z})$ acts on the upper halfspace model of the hyperbolic plane $\mathbb{H}^2$ by Möbius transformations, preserving the Farey triangulation. We have drawn the Farey triangulation below, in the disc model.



Figure 1: The Farey triangulation in the disc model of $\mathbb{H}^2$.

The quotient $SL(2,\mathbb{Z})\backslash\mathbb{H}^2$ is a hyperbolic orbifold with a cusp. It is often referred to as the modular surface, or as the $(2,3,\infty)$-triangle orbifold, as there is a fundamental domain for the action which is a triangle with one ideal vertex, and the other vertices of angles $\pi/2$ and $\pi/3$, and each ideal triangle in the Farey triangulation contains six $(2,3,\infty)$-triangles.

We now describe two different ways of choosing a point at infinity on the boundary. Given a basepoint x_0 in $\mathbb{H}^2$, we may identify the circle at infinity with the collection of geodesic rays based at x_0, which we may also identify with the unit tangent bundle. The unit tangent bundle has a natural measure arising from the Riemannian metric, and we will refer to this measure as Lebesgue measure.

Alternatively, we may choose a geodesic by taking a random walk on the group $SL(2,\mathbb{Z})$. By the Milnor-Švarc lemma, the Cayley graph of $SL(2,\mathbb{Z})$ is quasi-isometric to the infinite trivalent tree that is dual to the Farey triangulation. For simplicity, we assume that we are doing a simple random walk on this dual tree. We may identify points in the boundary of the tree with points on $S^1 = \partial\mathbb{H}^2$. A random walk on such a tree converges to the boundary almost surely, and this gives a hitting measure on S^1, which we shall call harmonic measure. We may then choose a geodesic from the basepoint x_0 to the chosen point at infinity.

The two measures on the boundary are in fact mutually singular, and furthermore, we can describe sets which have full measure in one measure, and measure zero in the other measure, in terms of the behaviour of the geodesics.



Figure 2: A geodesic in the Farey triangulation.

A geodesic from the basepoint to a particular number $r \in S^1$ passes through some sequence of fundamental domains, or equivalently, corresponds to a particular path in the trivalent tree converging to r at infinity. Starting from the basepoint, we may describe this path by a sequence of right and left turns, and these descriptions determine the continued fraction expansion of r. So if we write r as

$$r = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \cdots}}$$

then the a_i's correspond to the number of consecutive right and left turns along the path in the dual tree [Ser81]. Figure 2 shows a geodesic which starts off with three right turns followed by two left turns, so the geodesic will end on the circle at infinity somewhere between $1/3$ and $2/7$. It is a classical result, going back to Gauss, that for large i the proportion of numbers with continued fraction expansions containing $a_i = n$ is proportional to $1/n^2$ [Khi97]. If for simplicity we assume that the random walk on the tree has no back-tracking, then for harmonic measure, the left and right turns occur independently with equal probability, so the probability that $a_i = n$ is $1/2^n$. This means that "typical" geodesics for each measure have different distributions for the a_i. More precisely, we say that a geodesic has a $1/n^2$ distribution for the a_i, if the proportion of coefficients with $a_i = n$, for $0 \leqslant i \leqslant N$, is of the order of $1/n^2$, as $N \to \infty$. Similarly, we say a geodesic has an exponential distribution for the a_i if the proportion of geodesics with $a_i = n$, for $0 \leqslant i \leqslant N$, is of the order of $1/2^n$ as $N \to \infty$. These two sets of geodesics are sets which exhibit the singularity of the measures. The geodesics with a $1/n^2$-distribution have measure one with respect to Lebesgue measure, and measure zero with respect to harmonic measure. Similarly, the geodesics with an exponential distribution have measure zero with respect to Lebesgue measure, and measure one with respect to harmonic measure.

We wish to show an analogue of this result for Fuchsian groups, and the mapping class group. However, we find it convenient to interpret the distinct distributions above in terms of growth rates of the word metric along the geodesic. We now explain what this means in terms of $SL(2, \mathbb{Z})$. A point on a geodesic ray corresponds to a particular fundamental domain, which in turn corresponds to a particular (finite) sequence of left and right turns, $a_1, \ldots, a_n$. There is a group element g corresponding to the fundamental domain, and each left or right turn corresponds to adding a fixed number of generators to the word length, so the word length $d(1, g)$ of g is proportional to the sum of the a_i. However, there is an alternative metric d_{rel} we can use, which is the metric arising from the Farey graph by treating each edge as having length one. This metric is improper as the Farey graph is not locally finite, and this metric is often referred to as a relative metric on $SL(2, \mathbb{Z})$, as it is quasi-isometric to a word metric arising from an infinite generating set, consisting of a finite generating set, union a single parabolic subgroup. The distance in this metric is proportional to the number of changes from consecutive right turns to consecutive left turns, or vice versa, and so this is proportional to n. We may compare the two metrics on points along the geodesic, i.e. we may consider

$$\rho_n = \frac{d(1, g)}{d_{rel}(1, g)} \propto \frac{1}{n} \sum_{i=1}^{n} a_i.$$

In the case of harmonic measure, in which the coefficients a_i have an exponential distribution, this ratio above is bounded almost surely. This follows from the ergodic theorem, as an exponential distribution has finite first moment. However, in the case of Lebesgue measure, in which the coefficients have a $1/n^2$-distribution, then the ratio above tends to infinity almost surely, as the $1/n^2$-distribution does not have finite first moment. We may therefore use the set of geodesics for which ρ_n stays bounded, and the set of geodesics for which ρ_n tends to infinity, to exhibit the mutual singularity of harmonic measure and Lebesgue measure.

It is this last observation which we generalise to Fuchsian groups and the mapping class group. The word metric we use is the standard word metric with respect to a finite generating set for the group. The relative metric is a word metric with respect to an infinite generating set. In the Fuchsian case, this corresponds to adding a parabolic subgroup for each cusp to the generating set. In the case of the mapping class group we add subgroups which stabilize representatives of orbits of simple closed curves, and this is quasi-isometric to the orbit metric arising from the action of the mapping class group on the complex of curves.

1.2 Outline

We now give a brief outline of how we obtain the result for the mapping class group G acting on Teichmüller space $\mathcal{T}(S)$.

We shall consider Teichmüller space $\mathcal{T}(S)$ with the Teichmüller metric. There is a measure μ_{hol}, called holonomy measure or Masur-Veech measure, on $\mathcal{T}(S)$ for which the Teichmüller geodesic flow is ergodic, and for which the moduli space $\mathcal{M}(S) = G\backslash\mathcal{T}(S)$ has finite volume. The analogue of a cusp in hyperbolic space is the *thin part* $\mathcal{M}_\epsilon$, which consists of all points $X \in \mathcal{M}(S)$ which correspond to hyperbolic surfaces which have a curve of length at most ϵ. The pre-image of the cusp of a hyperbolic surface in the universal cover $\mathbb{H}^2$ is a disjoint union of horoballs. In Teichmüller space, the analogue of these horoballs is the *thin part* $\mathcal{T}_\epsilon$, which is the pre-image of $\mathcal{M}_\epsilon$ in $\mathcal{T}(S)$. However, this analogy is inexact, as $\mathcal{T}_\epsilon$ is connected in $\mathcal{T}(S)$, and there is no analogue of the disjointness property that the horoballs possess in hyperbolic space.

The cotangent bundle to Teichmüller space is the space of unit area quadratic differentials $\mathcal{Q}$. As a unit area quadratic differential corresponds to a flat surface, there is an action of $SL(2, \mathbb{R})$ on $\mathcal{Q}$, and the orbit of a point $q \in \mathcal{Q}$ gives an embedding of $SL(2, \mathbb{R})$ in $\mathcal{Q}$, which we shall denote $\widetilde{D}_q = SL(2, \mathbb{R}) \cdot q$.

In particular, the group of rotations of $\mathbb{R}^2$ acts on $\mathcal{Q}$, while the action is trivial on Teichmüller space. Indeed, a rotation by angle θ of the flat surface sends $(X, q) \mapsto (X, e^{-2i\theta}q)$. It follows from the definition that holonomy measure is invariant under rotation; as a consequence, the conditional measure from μ_{hol} on the circle $\{e^{i\theta}q : \theta \in [0, 2\pi]\}$ is precisely the Lebesgue (or Haar) measure.

A *metric cylinder* on a flat surface is a subset of the surface isometric to a flat cylinder, obtained by identifying two opposite sides of a parallelogram. The *length* of the metric cylinder is the length of a geodesic core curve in the flat metric. For a fixed metric cylinder, the subset of D_q corresponding to flat surfaces which contain metric cylinders of length at most L is a horoball in D_q. Masur [Mas93] showed that each Teichmüller disc contains many such disjoint horoballs corresponding to metric cylinders with area bounded below; in fact, sufficiently many that almost all geodesics from the basepoint spend a definite proportion of its time in the horoballs, This is known as a *good* packing of horoballs in Sullivan's terminology [Sul82]. As there is a lower bound on the area of the metric cylinder, we may find upper bounds for the length of the core curve of the cylinder in the hyperbolic metric, and so these horoballs are contained in the thin part $\mathcal{T}_\epsilon$.

When a geodesic γ passes through a horoball H, the distance it travels into the horoball is comparable to the amount of time it spends in the horoball. We say the *excursion* $E(\gamma, H)$ is the distance between the endpoints in the horoball, in the hyperbolic metric on the boundary of the horoball, and this is exponential in the amount of time the geodesic spends in the horoball (Section 2.6). By ergodicity of the Teichmüller geodesic flow, we show that the sum of the excursions grows faster than linearly along the geodesic (Theorem 3.9). Using results of Rafi, we show that the excursion is comparable to the twist parameter along the core curve of the corresponding metric cylinder, and that this can be used to estimate the change in the subsurface projection to the core curve along the geodesic. We may then estimate the growth of word length along the geodesic using the Masur-Minsky formula for word length in terms of subsurface projections, and we show that this in turn grows strictly faster than linearly along the geodesic. On the other hand, the relative metric grows at most linearly in time, hence the proof of the first part of Theorem 1.3 (Section 3.4).

The estimate for the hitting measure follows from the fact that along a random sample path both the word metric and the relative metric grow linearly in the number of steps [Mah10a], together with a sublinear estimate of the deviation between geodesic and sample path along the lines of [Tio12] (Section 4).

Finally, in Section 5 we show how to apply this method to the simpler case of Fuchsian groups, thus proving Theorem 1.1.

1.3 Acknowledgements

We would like to thank Howard Masur, Curtis McMullen, Kasra Rafi and John Smillie for helpful conversations.

2 Preliminaries

In this section we review some background and previous results that we use, and set up some notation.

2.1 Notation

We shall find it convenient to occasionally use *big O* notation. We say that $f(x) = O(g(x))$ if there are constants A and B such that $|f(x)| \leqslant A\,|g(x)|$ for all $x \geqslant B$. In particular, $f(x) = O(1)$ means that the function $f(x)$ is bounded. We will also write $f(x) \asymp g(x)$ to mean that $f(x)$ and $g(x)$ are equivalent up to additive and multiplicative constants, i.e. there are constants K and c such that

$$\frac{1}{K} g(x) - c \leqslant f(x) \leqslant K g(x) + c.$$

2.2 Quadratic differentials and Teichmüller discs

Let S be a hyperbolic surface of finite type, i.e. a surface of finite area which may have boundary components or punctures. We say such a surface S is *sporadic* if it is a sphere with at most four punctures or boundary components, or a torus with at most one puncture or boundary component. We shall primarily be interested in non-sporadic surfaces.

Let S be a non-sporadic surface with no boundary components, but which may have punctures. We will write $\mathcal{T}(S)$ for the Teichmüller space of a surface S, or just $\mathcal{T}$ if we do not need to explicitly refer to the surface. We shall consider $\mathcal{T}$ together with the Teichmüller metric

$$d_{\mathcal{T}}(x, y) = \tfrac{1}{2} \inf_{f} \log K(f),$$

where the infimum is taken over all quasiconformal maps $f \colon x \to y$, and $K(f)$ is the quasiconformal constant for the map f. The mapping class group $G = \text{Mod}(S)$ of the surface acts by isometries on $\mathcal{T}$, and we shall write $\mathcal{T}_\epsilon$ for the *thin part* of Teichmüller space, i.e. all surfaces which contain a curve of hyperbolic length at most ϵ. We shall write $\mathcal{M}$ for the quotient $G \backslash \mathcal{T}$, which is known as moduli space. The thin part of Teichmüller space is mapping class group invariant, and we shall write $\mathcal{M}_\epsilon$ for the subset of moduli space given by $G \backslash \mathcal{T}_\epsilon$.

Let $\mathcal{Q}$ be the space of unit area quadratic differentials, which may be identified with the unit cotangent bundle to Teichmüller space. We shall write π for the projection $\pi : \mathcal{Q} \to \mathcal{T}$ which sends a quadratic differential to its underlying Riemann surface, and we shall write μ_{hol} for the Masur-Veech measure, also known as the holonomy measure, as it may be defined in terms of holonomy coordinates. The measure μ_{hol} is mapping class group invariant, and so gives a measure on the moduli space of unit area quadratic differentials $\mathcal{MQ} = G \backslash \mathcal{Q}$, which has finite volume.

A quadratic differential q determines a flat structure on the surface, which may be thought of as a union of polygons glued together along parallel sides, where the vertices of the polygons may correspond to points of cone angle $n\pi$, for $n \geqslant 1$. If $n \geqslant 2$, then the vertex corresponds to a zero of order $n-2$ for the quadratic differential q, and for $n = 1$ the vertices correspond to cone points of angle π which are simple poles for the quadratic differential, and correspond to the punctures of the surface. There is an affine action of $SL(2, \mathbb{R})$ on the flat surface, which gives rise to a new quadratic differential. The orbits of quadratic differentials under the action of $SL(2, \mathbb{R})$ give a foliation of $\mathcal{Q}$ by copies of $SL(2, \mathbb{R})$, and we shall write $\widetilde{D}_q$ for the orbit of the quadratic differential q. We shall write D_q for the image of $\widetilde{D}_q$ in $\mathcal{T}$, and this is called a Teichmüller disc, which is geodesically embedded in $\mathcal{T}$. With the metric induced from the Teichmüller metric, D_q is isometric to the hyperbolic plane of constant curvature -4, and it will be convenient for us to use coordinates coming from the disc model of hyperbolic plane, with the initial quadratic differential q corresponding to the origin.

The group of rotations of $\mathbb{R}^2$ acts on flat surfaces, and hence on $\mathcal{Q}$. In terms of quadratic differentials, rotation by angle θ in $\mathbb{R}^2$ sends $q \mapsto e^{-2i\theta} q$, and this action is trivial on Teichmüller space $\mathcal{T}$. It follows from the definition that holonomy measure is invariant under rotation, i.e. $\mu_{\text{hol}}(U) = \mu_{\text{hol}}(e^{i\theta} U)$, for all θ, for any subset $U \subset \mathcal{Q}$. In particular, this means that if we consider the conditional measure from μ_{hol} on the image of a point $q \in \mathcal{Q}$ under rotation, i.e. $\{e^{i\theta} q : \theta \in [0, 2\pi]\}$, then this is precisely the invariant Haar or Lebesgue measure on the circle.

2.3 Curve complex and subsurface projections

In this section we review the properties we will use of two combinatorial objects associated with a surface, namely the *curve complex* and the *marking complex.*

We say a simple closed curve on a surface S is *essential* if it does not bound a disc, and is not parallel to a puncture or boundary component. The *curve complex* $\mathcal{C}(S)$ is a finite dimensional but locally infinite simplicial complex whose vertices are isotopy classes of essential simple closed curves on S, and whose simplices consist of collections of curves which can be realised disjointly on the surface. For the non-sporadic surfaces, the curve complex is a non-empty connected simplicial complex. In the case of a torus with one puncture or boundary component, or a sphere with four punctures or boundary components, the definition above gives a complex with no edges, so we alter the definition to connect two vertices if their corresponding curves can be realised by curves which intersect at most once (in the case of the once punctured torus) or at most twice (in the case of the four punctured sphere). In the case of the annulus the curve complex is defined to be the infinite graph with vertices consisting of arcs connecting the two boundary components of the annulus modulo isotopy fixing the endpoints with edges between two arcs if they can be realized disjointly. The curve complex of the annulus is quasi-isometric to $\mathbb{Z}$ with a quasi-isometry given by the algebraic intersection number. We define the curve complex to be empty for the remaining sporadic surfaces.

We say a subsurface $Y \subseteq S$ is *essential* if each boundary component is an essential simple closed curve in S. Given an essential subsurface $Y \subseteq S$, which is not a disc or a three-punctured sphere, one can also consider $\mathcal{C}(Y)$, the complex of curves of Y. There is a coarsely well-defined *subsurface projection* $\pi_Y : \mathcal{C}(S) \to \mathcal{C}(Y)$ which we now describe. Choose an element in the isotopy class of the curve γ which has the minimal possible number of intersections with Y, and then take a regular neighbourhood of the union of the boundary of Y with the intersection of the curve γ with Y, i.e. $N(\partial Y \cup (\gamma \cap Y))$. Choose a component of the boundary of this regular neighbourhood to be $\pi_Y(\gamma)$. This is coarsely well defined.

To define the annular projection $\pi(\gamma)$ of a curve γ with essential intersection with an annulus A essentially one passes to the annulus cover $\widetilde{S}$ of S given by the core curve α of A and chooses $\pi_A(\gamma)$ to be a component of the lift of γ that is an arc running from one boundary component of $\widetilde{S}$ to the other. The set of components of the lift of γ that satisfy this property form a finite diameter set in the curve complex of the annulus $\widetilde{S}$ and so the projection is coarsely well-defined. Finally the map π_A has the property that if D_α denotes the Dehn twist about α, then

$$d_{\mathcal{C}(A)}(\pi_A(D_\alpha^n(\gamma)), \pi_A(\gamma)) = 2 + |n|. \tag{1}$$

Thus, defining the projection this way achieves the desired property of recording the twisting around α. There is a natural $\mathbb{Z}$ action on $\mathcal{C}(A)$ by Dehn twisting around the core curve of the annular cover $\widehat{S}$. The group $\mathbb{Z}$ also has an inclusion into the mapping class group of S as Dehn twists around α, and so it acts on $\mathcal{C}(A)$ through this inclusion. The projection map π_A is coarsely equivariant with respect to the two $\mathbb{Z}$ actions. we will often abuse notation and write π_α to mean the subsurface projection to an annulus whose core curve is α.

A *marking* consists of a collection of simple closed curves α_i forming a maximal simplex in the curve complex, or equivalently, a pants decomposition of the surface, together with a *transverse curve* τ_i for each pants curve α_i, which is an element of the annular curve complex corresponding to α_i. The curves α_i are known as the *base curves* of the marking. We remark that the definition we give here corresponds to the definition of a *complete* marking from [MM00]. They consider more general markings, in which the set of base curves does not need to form a maximal simplex in $\mathcal{C}(S)$, and all base curves are not required to have a transversal. However, complete markings suffice for our purposes.

If α is a simple closed curve in S, then a *clean transverse curve* for α is a simple closed curve β, such that a regular neighbourhood of $\alpha \cup \beta$, isotoped to have minimal intersection, is either a sphere with four boundary components, or a torus with a single boundary component. A *clean marking* is a marking (α_i, τ_i), such that each transverse curve τ_i is of the form $\pi_{\alpha_i}(\beta_i)$, for some clean transverse curve β_i, which is disjoint from the union of the other base base curves $\cup \alpha_j$, for $j \neq i$. A clean marking $m' = (\alpha_i, \beta_i)$ is compatible with a marking $m = (\alpha_i, \tau_i)$, if the base of m is equal to the base of m', and for each simple closed curve α_i in the base, $d_{\alpha_i}(\tau_i, \pi_{\alpha_i}\beta_i)$ is minimal. There are only finitely many clean markings m' compatible with a given marking m.

The *marking complex* $M(S)$ is a graph whose points are clean markings, and whose edges are given by *elementary moves* as defined by Masur and Minsky [MM00]. These moves are called twists and flips. In a *twist*, a transverse curve β_i is replaced by the image of the transverse curve under a Dehn twist along its corresponding pants curve $D_{\alpha_i}(\beta_i)$. In a *flip*, a transverse curve β_i and its corresponding base curve α_i are interchanged, i.e. a new clean marking is chosen which is compatible with the marking formed by replacing $(\alpha_i, \pi_{\alpha_i}(\beta_i))$ with $(\beta_i, \pi_{\beta_i}(\alpha))$. The mapping class group acts on the marking complex and the space of orbits is finite. We will write d_M for the induced metric on the marking complex obtained by setting the length of each edge equal to one.

The mapping class group is finitely generated, so a choice of generating set gives rise to a word metric, in which the length of a group element is the shortest length of any product of generators representing the group element. Different generating sets give rise to quasi-isometric metrics. We shall assumed we have fixed a generating set, and we shall write d_G for the word metric distance in the mapping class group. Masur and Minsky showed that the distance d_M in the marking complex is quasi-isometric to the word metric d_G in the mapping class group.

Proposition 2.1. [MM00, Theorems 6.10 and 7.1] *Fix a complete clean marking m_0 and a system of generators for* $\text{Mod}(S)$. *Then there exist constants C_1, C_2 such that for each $g \in \text{Mod}(S)$*

$$C_1^{-1} d_G(1, g) - C_2 \leqslant d_M(m_0, gm_0) \leqslant C_1 d_G(1, g) + C_2$$

There is a coarsely well-defined map from the marking complex $M(S)$ to the curve complex $\mathcal{C}(S)$, which takes a marking to one of the short curves in the marking. In particular, for any essential subsurface $Y \subseteq S$, this gives us a map from the marking space $M(S)$ to $\mathcal{C}(Y)$, given by composing π and π_Y. Given markings m and n, denote by $d_Y(m, n)$ the diameter in $\mathcal{C}(Y)$ of the union of the projections of m and n. If α is a simple closed curve, then d_α will denote the distance in the curve complex of the annulus with core curve α.

Masur and Minsky [MM00, Theorem 6.12] proved a distance formula expressing the distance in the marking complex $M(S)$, and hence by Proposition 2.1, the distance in $\text{Mod}(S)$ in the word metric, in terms of subsurface projections. Here, we state a slightly simplified version of it. Given a number $A > 0$, for any number d, we define a cutoff function given by

$$\lfloor d \rfloor_A = \begin{cases} d & \text{if } d \geqslant A \\ 0 & \text{otherwise.} \end{cases}$$

Theorem 2.2 ([MM00] Quasi-distance formula)**.** *There exists a constant $A_0 > 0$, which depends only on the topology of the surface S, such that for any $A \geqslant A_0$, there are constants C_1 and C_2, which depend only on A and the topology of S, such that for any pair of clean markings m and m' in $M(S)$,*

$$C_1^{-1} d_M(m, m') - C_2 \leqslant \sum_{Y \subseteq S} \lfloor d_Y(m, m') \rfloor_A \leqslant C_1 d_M(m, m') + C_2$$

where the sum runs over all subsurfaces Y of S, including S.

2.4 Short curves and short markings

Given a hyperbolic surface X, there is a *systole* map from Teichmüller space $\mathcal{T}(S)$ to the curve complex $\mathcal{C}(S)$ given by sending X to a shortest curve on X. This map is coarsely well defined: there may be multiple shortest curves, but there are only finitely many choices, and they are a bounded distance apart in the curve complex, where these bounds depend only on the topology of S. This follows from the fact that by Bers' Lemma, for any surface S there is a constant L depending on S such that any hyperbolic metric on S contains a simple closed curve of length at most

L, and the collar lemma says that for any simple closed curve γ of length L, there is an $\epsilon > 0$, depending on L, such that an ϵ-neighbourhood of γ is embedded, and so this bounds the number of intersections of any pair of curves of length L. In particular, for any Teichmüller geodesic γ_t, this gives a sequence of simple closed curves α_t.

A *reparameterization* of $\mathbb{R}$ is a continuous, monotonically increasing function $\phi\colon \mathbb{R} \to \mathbb{R}$, which need not be onto. We say a function f from $\mathbb{R}$ to a metric space is an *unparameterized* (K, c)*-quasigeodesic* if there is a reparameterization ϕ such that $f \circ \phi$ is a (K, c)-quasigeodesic, which may be of finite length.

Masur and Minsky [MM00] showed that the image of a Teichmüller geodesic under the shortest curve map is an unparameterized quasigeodesic in the curve complex. Rafi [Raf10] showed that the composition of subsurface projection with the shortest curve map gives an unparameterized quasigeodesic in the curve complex of the subsurface.

Theorem 2.3 ([Raf10, Theorem B]). *There are constants K and c, which only depend on the surface S, such that for any Teichmüller geodesic γ, and any subsurface $Y \subseteq S$, the sequence of curves $\pi_Y(\alpha_t)$ arising from the projection of the shortest curves α_t to $\mathcal{C}(Y)$ is an unparameterized (K, c)-quasigeodesic in $\mathcal{C}(Y)$.*

Given a hyperbolic surface X, let us define the *shortest marking* $m(X)$ in the following way. First, choose a pants decomposition by picking the shortest simple closed curves in the hyperbolic metric, using the greedy algorithm. To be precise, start by choosing one of the shortest curves on the surfaces, then choose one of the shortest curves on the complementary surface, and continue until you have a pants decomposition of the original surface. Then, for each curve α_i of the pants decomposition choose a transverse curve τ_i which is perpendicular to α_i in the hyperbolic metric. If there are multiple shortest curves, then the shortest marking may not be unique, but there are only a finite number of choices, with a bound depending on the topology of the surface. This gives a map from $\mathcal{T}(S)$ to $M(S)$, which is coarsely well-defined, and we shall write m_t for the image of a point on a Teichmüller geodesic γ_t under this map.

2.5 Projections and isolated intervals

Rafi [Raf07] shows that for any Teichmüller geodesic γ, and any subsurface Y, there is a (possibly empty) interval I_Y during which Y is *isolated*, i.e. the boundary components of Y are short in the hyperbolic metric. In order to make this statement precise, let us pick a constant $\epsilon_0 > 0$ which is smaller than the Margulis constant. Given a Teichmüller geodesic $\gamma(t)$, and a simple closed curve α, we shall write $L_t(\alpha)$ for the length of α in the hyperbolic metric $\gamma(t)$.

Proposition 2.4 ([Raf07, Corollary 3.3]). *Let $\epsilon_0 > 0$ be sufficiently small. Then there exists $\epsilon_1 \leqslant \epsilon_0$ such that, for any geodesic in the Teichmüller space and any curve α in S, there exists a connected (perhaps empty) interval I_α such that*

1. *For $t \in I_\alpha$, $L_t(\alpha) \leqslant \epsilon_0$*
2. *For $t \notin I_\alpha$, $L_t(\alpha) \geqslant \epsilon_1$*

Outside the active interval I_α, the map from the Teichmüller geodesic to the curve complex of the annulus corresponding to α is coarsely constant.

Proposition 2.5 ([Raf07, Proposition 3.7])**.** *There is a constant K, depending only on the topology of the surface S, and the choices for the constants ϵ_0 and ϵ_1, such that if $[r,s] \cap I_\alpha = \emptyset$, then*

$$d_\alpha(m_r, m_s) \leqslant K.$$

In the next section we show that the length of the active interval for an annulus is roughly log of the projection distance of the endpoints of the geodesic into the subsurface.

2.6 Excursions and twist parameter

The material in this section is due to Rafi [Raf07, Raf10]. However, we need versions of his results in terms of the excursion parameter, and we use some of the contents of the proofs, not just the main stated results, so we write out all of the details for the convenience of the reader.

A horoball H in the hyperbolic plane is a subset of the plane which in the Poincaré disc model corresponds to a Euclidean disc whose boundary circle is tangent to the boundary at infinity. Given a horoball H and a geodesic γ which spends a finite amount of time in H, let us define the *excursion* $E(\gamma, H)$ of γ in H as the "relative visual size" of the set of rays which go deeper than γ inside H. Namely, consider a basepoint X_0 on the Teichmüller disc in $\mathcal{T}$, and let γ_H be the geodesic through X_0 which tends to the cusp of H, and γ_T a geodesic through X_0 which is tangent to H. Let ϕ_0 be the angle between γ and γ_H, and ϕ_{max} be the angle between γ_H and γ_T (see Figure 3). Then

Definition 2.6.

$$E(\gamma, H) := \frac{\phi_{max}}{\phi_0} \tag{2}$$

It turns out that $E(\gamma, H)$ is, up to an additive error, also the hyperbolic length of the projection of $\gamma \cap H$ to the complement of H.



Figure 3: Excursion in the horoball H.

Let (X, q) be a quadratic differential on X, and α a simple closed curve on X. The choice of q determines a Teichmüller geodesic γ and a pair (F^+, F^-) of contracting and expanding foliations. Each t determines a new quadratic differential q_t and hence a flat metric on X, which we will call the q_t-metric.

For a given t, α is realized by a family of parallel flat geodesics, and we will denote as β_t the perpendicular to α in the q_t-metric. The *twist parameter* $tw_t^+(\alpha)$ is the highest intersection number between a leaf of F^+ and the transversal β_t, and similarly we define $tw_t^-(\alpha)$.

Given a simple closed curve α corresponding to metric cylinder, there is a unique rotation $e^{i\theta_\alpha}$ which takes the metric cylinder to a vertical metric cylinder. The endpoint of the geodesic ray corresponding to the quadratic differential $e^{i\theta_\alpha}q$ determines a point ξ_α on the boundary at infinity of the Teichmüller disc $\mathbb{D}$. We shall write $H_\epsilon(\alpha)$ as the set of points in the disc for which α is short in the flat metric:

$$H_\epsilon(\alpha) := \{q \in \mathbb{D} \ : \ell_q^2(\alpha) \leqslant \epsilon\}$$

As seen in the disc, this set is a horoball the tangent to the boundary at infinity at ξ_α. The fundamental estimate is the following:

Proposition 2.7. *Let $H = H_\epsilon(\alpha)$ as above, and let t_1 and t_2 respectively be the entry time and exit time from H (i.e. $t_1 \leqslant t_2$) along the Teichmüller geodesic γ. Let moreover A be the area of the maximal flat cylinder in (X, q_0) with core curve α. Then*

$$tw_{t_2}^-(\alpha) - tw_{t_1}^-(\alpha) \asymp \frac{A}{\epsilon} E(\gamma, H)$$

Proof. Consider the universal cover of the flat cylinder corresponding to α at time t, in the flat metric q_t. We shall assume that the contracting foliation is vertical, and the expanding foliation is horizontal. Let ℓ_t be the length of α at time t, and let θ_t be the angle α_t makes with the vertical contracting foliation, as illustrated below in Figure 4.



Figure 4: Estimating intersections in the flat annulus.

Let h_t and v_t be the horizontal and vertical lengths of ℓ_t in the q_t metric, i.e.

$$h_t = h_0 e^t = \ell_0 \sin\theta_0 e^t$$
$$v_t = v_0 e^{-t} = \ell_0 \cos\theta_0 e^{-t}$$

Let w_t be the length of β_t, which is the width of the flat annulus. Let H_t be the length of the intersection of a leaf of the horizontal foliation with the universal cover of the flat annulus, and let H'_t be the length of the intersection of the horizontal leaf with two adjacent translates of β_t.

Up to constant additive error, $tw_t^-(\alpha)$, which is the maximum number of intersections between the horizontal leaf of the foliation and β, is given by H_t/H'_t. Therefore

$$tw_t^-(\alpha) = \frac{H_t}{H'_t} + O(1) = \frac{w_t \sin\theta_t}{\ell_t \cos\theta_t} + O(1).$$

The area of the annulus is $A = w_t \ell_t$, and $\tan\theta_t = \tan\theta_0 e^{2t}$, so this implies that

$$tw_t^-(\alpha) = \frac{A}{\ell_t^2} \tan\theta_0 e^{2t} + O(1). \tag{3}$$

The total length of α is given by

$$\ell_t^2 = h_t^2 + v_t^2 = \ell_0^2(\sin^2\theta_0 e^{2t} + \cos^2\theta_0 e^{-2t}), \tag{4}$$

and recall that we choose t_i such that $\ell_{t_i}^2 = \epsilon$, which by (3) implies

$$tw_{t_2}^-(\alpha) - tw_{t_1}^-(\alpha) = \frac{A}{\epsilon} \tan\theta_0 (e^{2t_2} - e^{2t_1}) + O(1) \tag{5}$$

Note that by definition the t_i are solutions to the equation

$$\ell_{t_i}^2 = \ell_0^2(\sin^2\theta_0 e^{2t_i} + \cos^2\theta_0 e^{-2t_i}) = \epsilon \qquad i = 1, 2$$

If we set $X_i := e^{2t_i}$, then X_i are the solutions to

$$X^2 - \frac{\epsilon^2}{\ell_0^2 \sin^2\theta_0} X + \frac{1}{\tan^2\theta_0} = 0 \tag{6}$$

hence

$$e^{2t_2} - e^{2t_1} = X_2 - X_1 = \sqrt{\frac{\epsilon^2}{\ell_0^4 \sin^4\theta_0} - \frac{4}{\tan^2\theta_0}} \tag{7}$$

and putting (6) and (7) together

$$tw_{t_2}^-(\alpha) - tw_{t_1}^-(\alpha) = \frac{A\tan\theta_0}{\epsilon}(e^{2t_2} - e^{2t_1}) + O(1) = \frac{A}{\epsilon}\sqrt{\frac{\epsilon^2}{\ell_0^4 \sin^2\theta_0 \cos^2\theta_0} - 4} + O(1) \tag{8}$$

Let us now relate this quantity to the excursion in the horoball H.

Let ϕ_{max} be the angle between a geodesic tangent to H and the geodesic γ_H which goes straight into the cusp of H. We have the

Lemma 2.8.

$$\sin\phi_{max} = \frac{\epsilon}{\ell_0^2}$$

Proof. When $\theta_0 = \theta_{max}$ then the geodesic is tangent to the horoball H, hence $t_1 = t_2$ in eq. (7), so

$$\frac{\epsilon^2}{\ell_0^4 \sin^4\theta_{max}} = \frac{4}{\tan^2\theta_{max}}$$

The claim follows by recalling that a rotation of angle θ in the flat metric picture corresponds to multiplying the quadratic differential by $e^{2i\theta}$, hence $\phi_{max} = 2\theta_{max}$. □

The proposition now follows easily from the lemma, eq. (8) and the fact that $\phi_0 = 2\theta_0$:

$$tw^-_{t_2}(\alpha) - tw^-_{t_1}(\alpha) = \frac{2A}{\epsilon}\frac{\sin\phi_{max}}{\sin\phi_0}\sqrt{1 - \frac{\sin^2\phi_0}{\sin^2\phi_{max}}} + O(1) \asymp \frac{A}{\epsilon}E(\gamma, H)$$

where in the last equality we used equation (2) and the fact that $\sin\phi \asymp \phi$. $\square$

Remark. *Note that one can also relate the twist parameter to the time spent by the geodesic inside the horoball, namely*

$$tw^-_{t_2}(\alpha) - tw^-_{t_1}(\alpha) \asymp \frac{A}{\epsilon}(e^{t_2-t_1} - e^{t_1-t_2})$$

The distance between the projections from the marking complex to the complex of the annulus can be compared to the excursion in the horoball:

Proposition 2.9. *Let $\epsilon > 0$ sufficiently small, and (X_0, q) a unit area quadratic differential, which determines the geodesic ray γ_t. Let A be the q-area of the maximal flat cylinder with core curve α, and suppose that α is not short in the q-metric (i.e. $\ell_q^2(\alpha) \geq \epsilon$). If the geodesic γ crosses the horoball $H = H_\epsilon(\alpha)$ and t is larger than the exit time of γ from H, then*

$$d_\alpha(m_0, m_t) \asymp \frac{A}{\epsilon}E(\gamma, H)$$

where m_t is the shortest marking on γ_t, and the quasi-isometry constants depend only on X_0, ϵ and the topology of S.

Before proving the proposition, let us recall the definition of extremal length:

$$\mathrm{Ext}_\sigma(\alpha) := \sup_\rho \frac{(\ell_\rho(\alpha))^2}{A(\rho)}$$

where the sup is taken over all metrics ρ in the same conformal class as σ. For any quadratic differential q with area 1 and any curve α,

$$(\ell_q(\alpha))^2 \leqslant \mathrm{Ext}_\sigma(\alpha) \leqslant \frac{L_\sigma(\alpha)}{2}e^{L_\sigma(\alpha)/2}$$

where the left-hand side is by definition, while the right-hand side is due to Maskit [Mas85], and $L_\sigma(\alpha)$ is the length of α in the hyperbolic metric corresponding to the conformal structure σ.

Recall $tw_q^\pm$ denotes the twist parameter in the flat metric associated to q, as defined in the previous section. Analogously, given a hyperbolic metric σ on S and a simple closed curve α, we can define a twist parameter $tw_\sigma^\pm(\alpha)$ with respect to the hyperbolic metric by taking a curve β perpendicular to α with respect to the hyperbolic metric and letting

$$tw_\sigma^\pm(\alpha) := i(F^\pm, \beta)$$

The following proposition of Rafi relates the two twist parameters:

Proposition 2.10 ([Raf07, Theorem 4.3])**.** *The two twist parameters are the same up to an additive error comparable to $1/L_\sigma(\alpha)$. That is,*

$$tw_\sigma^\pm(\alpha) = tw_q^\pm(\alpha) + O\left(\frac{1}{L_\sigma(\alpha)}\right)$$

Proof of Proposition 2.9. Let us choose ϵ_0 in such a way that if $\ell_q^2(\alpha) = \epsilon$, then $L_\sigma(\alpha) \geqslant \epsilon_0$. Let t_1 and t_2 be the times when the q_t-length of α is exactly ϵ, and let $t > t_2$. By the previous choice, $L_{\sigma_{t_i}}(\alpha) \geqslant \epsilon_0$ for $i = 1, 2$, so by Proposition 2.4 $[0, t_1]$ and $[t_2, t]$ are disjoint from I_α, hence by Proposition 2.5

$$d_\alpha(m_0, m_t) = d_\alpha(m_{t_1}, m_{t_2}) + O(1)$$

On the other hand, the progress in subsurface projection across the horoball is comparable to the progress in the twist parameter for the hyperbolic metric,

$$d_\alpha(m_{t_1}, m_{t_2}) = |i_\alpha(\beta_{t_1}, F^+) - i_\alpha(\beta_{t_2}, F^+)| + O(1) = |tw^+_{\sigma_1}(\alpha) - tw^+_{\sigma_2}(\alpha)| + O(1) =$$

and by Proposition 2.10 it is also comparable to the progress in the twist parameter defined via the flat metric:

$$= |tw^+_{q_1}(\alpha) - tw^+_{q_2}(\alpha)| + O\left(\frac{1}{L_{\sigma_1}(\alpha)}\right) + O\left(\frac{1}{L_{\sigma_2}(\alpha)}\right)$$

and since $L_{\sigma_i}(\alpha) \geqslant \epsilon_0$

$$d_\alpha(m_0, m_t) = |tw^+_{q_1}(\alpha) - tw^+_{q_2}(\alpha)| + O(1)$$

Finally, by Proposition 2.7 the twist is comparable to the excursion, thus

$$d_\alpha(m_0, m_t) \asymp \frac{A}{\epsilon} E(\gamma, H)$$

□

2.7 Coarse monotonicity for the word metric

In [Raf10], Rafi shows the following non-backtracking or reverse triangle inequality for subsurface projections along a Teichmüller geodesic. Recall that given a Teichmüller geodesic γ_t we write m_t for the shortest marking at γ_t, and we write $d_Y(m_s, m_t)$ to mean the distance in the curve complex $\mathcal{C}(Y)$ between the images of m_s and m_t under subsurface projection to Y.

Theorem 2.11 ([Raf10, Theorem 6.1]). *There exists a constant C, only depending on the topology of S, such that for every Teichmüller geodesic γ, and every subsurface Y,*

$$d_Y(m_r, m_s) + d_Y(m_s, m_t) \leqslant d_Y(m_r, m_t) + C, \tag{9}$$

for all constants $r \leqslant s \leqslant t$.

The above theorem along with the Masur-Minsky quasi-distance formula 2.2 implies that the distance in the marking complex is coarsely monotonic along a Teichmüller ray.

Proposition 2.12. *There exists constants $C_1 > 0$ and C_2 that depend only on S such that along a Teichmüller geodesic γ_t, for $0 < s < t$ the distance in the marking complex satisfies*

$$d_M(m_0, m_s) \leqslant C_1 d_M(m_0, m_t) + C_2.$$

Proof. Let C be the constant in Rafi's reverse triangle inequality, Theorem 2.11. Assume $0 < s < t$, then (9) implies

$$d_Y(m_0, m_t) \geqslant d_Y(m_0, m_s) - C, \tag{10}$$

for all subsurfaces $Y \subseteq S$. The Masur-Minsky quasi-distance formula (Theorem 2.2) holds for all floor constants sufficiently large, though the quasi-isometry constants depend on A. Choose a floor constant $A > 2C$, and let K_1 and K_2 be the associated quasi-isometry constants. By the definition of the floor function, if $\lfloor x \rfloor_A$ is non zero, then $x \geqslant A$. This implies that $x - A/2 \geqslant x/2$, and as the floor function is monotonic,

$$\lfloor x - A/2 \rfloor_A \geqslant \lfloor x/2 \rfloor_A. \tag{11}$$

As we have chosen $A > 2C$, combining (10) and (11) implies

$$\lfloor d_Y(m_0, m_t) \rfloor_A \geqslant \lfloor \tfrac{1}{2} d_Y(m_0, m_s) \rfloor_A, \tag{12}$$

again for all subsurfaces $Y \subseteq S$. Now summing (12) over all subsurfaces $Y \subseteq S$, the quasi-distance formula implies

$$d_M(m_0, m_t) \geqslant \frac{1}{K_1} \left(\sum \lfloor \tfrac{1}{2} d_Y(m_0, m_s) \rfloor_A - K_2 \right).$$

By definition of the floor function, $\lfloor \frac{1}{2} x \rfloor_A = \frac{1}{2} \lfloor x \rfloor_{2A}$, so

$$d_M(m_0, m_t) \geqslant \frac{1}{2K_1} \left(\sum \lfloor d_Y(m_0, m_s) \rfloor_{2A} - 2K_2 \right).$$

The quasi-distance formula holds for all A sufficiently large, so in particular holds for $2A$, though with different quasi-isometry constants, which we shall denote K_3 and K_4. This implies that

$$d_M(m_0, m_t) \geqslant \frac{1}{2K_1K_3} \left(d_M(m_0, m_s) - K_3K_4 - K_2 \right).$$

whence the result. ☐

2.8 Projection to closest Teichmüller lattice point

Let q be a quadratic differential, let q_t be the image of q under the Teichmüller geodesic flow after time t, and let X_t be the image of q_t in $\mathcal{T}$. The orbit of X_0 under the mapping class group is called a *Teichmüller lattice*, and let $h_t X_0$ be a choice of closest lattice point in $\mathcal{T}$ to X_t, i.e. such that

$$d_{\mathcal{T}}(h_t X_0, X_t) \leqslant d_{\mathcal{T}}(h X_0, X_t) \text{ for all } h \in \mathrm{Mod}(S).$$

For any given point X_t, there are at most finitely many closest lattice points, however it is possible that the number of closest lattice points increases as you choose points deeper in the thin part. Let m_t be a shortest marking on X_t, and d_G the word metric on the mapping class group with respect to some choice of generators.

Lemma 2.13. *If X_0 and X_t both lie in the thick part $\mathcal{T} \setminus \mathcal{T}_\epsilon$, then*

$$d_G(1, h_t) \asymp d_M(m_0, m_t)$$

where the quasi-isometry constants only depend on X_0, the choice of ϵ and the generating set for the mapping class group.

Proof. Let K_1 be the diameter of the thick part $\mathcal{T} \setminus \mathcal{T}_\epsilon$ in moduli space; then, by definition there exists a group element g such that in Teichmüller space $d_{\mathcal{T}}(gX_0, X_t) \leqslant K_1$, so by definition of h_t

$$d_{\mathcal{T}}(h_t X_0, X_t) \leqslant K_1.$$

Hence by group invariance

$$d_{\mathcal{T}}(X_0, h_t^{-1} X_t) \leqslant K_1.$$

In the Teichmüller ball of radius K_1 only finitely many markings appear as short markings, hence there exists K_2, depending only on K_1, and the surface S, such that the distance in the marking complex is bounded:

$$d_M(m_0, h_t^{-1} m_t) \leqslant K_2.$$

As a consequence,

$$|d_M(m_0, m_t) - d_M(m_0, h_t m_0)| \leqslant d_M(h_t m_0, m_t) = d_M(m_0, h_t^{-1} m_t) \leqslant K_2,$$

Finally, the distance in the word metric $d_G(1, h_t)$ is quasi-isometric to the distance $d_M(m_0, h_t m_0)$ in the marking complex by Proposition 2.1. □

By combining the previous lemma with the coarse monotonicity statement of Proposition 2.12, we get that the word length of the closest point projection to the Teichmüller lattice is coarsely monotone along the thick part of a Teichmüller ray:

Proposition 2.14. *There exists constants $C_1 > 0$ and C_2, that depend only on X_0 and ϵ_0 and the choice of generators, such that along a Teichmüller geodesic γ_t, for $0 < s < t$ the word metric satisfies*

$$d_G(1, h_s) \leqslant C_1 d_G(1, h_t) + C_2.$$

whenever γ_0, γ_s and γ_t all lie in the thick part $\mathcal{T} \setminus \mathcal{T}_{\epsilon_0}$.

3 Lebesgue measure sampling

The goal of this section is to study the asymptotic behaviour of typical Teichmüller geodesics with respect to Lebesgue measure, proving Theorem 1.3. More precisely, we want to keep track of short curves in the flat metric as the metric changes under the action of Teichmüller flow, and prove an asymptotic result (Theorem 3.3).

3.1 Metric cylinders with bounded area

Let q be a quadratic differential of unit area. A *metric cylinder* for q is a cylinder in the flat metric associated to q which is the union of freely homotopic closed trajectories of q. We shall label each metric cylinder by the homotopy class α of the corresponding closed trajectory.

Let us now fix some $0 < \delta < 1$, and let $C_q(\delta)$ be the set of metric cylinders for the q-metric with area bounded below by δ. Moreover, let us denote by $C_q(\delta, \epsilon)$ the set of cylinders whose area is bounded below by δ and whose core curve has length shorter than the square root of ϵ:

$$C_q(\delta, \epsilon) := \{\alpha \in C_q(\delta) \ : \ \ell_q^2(\alpha) \leqslant \epsilon\}$$

Lemma 3.1. *Suppose $\epsilon < \delta$. Then any two distinct elements of $C_q(\delta, \epsilon)$ are disjoint on q. As a corollary, the cardinality of $C_q(\delta, \epsilon)$ is bounded above by a constant which depends only on the topology of S.*

Proof. As in [Mas93], Lemma 2.2. Denote by α the core curve of some cylinder which belongs to $C_q(\delta, \epsilon)$. Since the metric cylinder of α has area $A(\alpha) \geqslant \delta$, any curve τ which crosses α is such that $\delta \leqslant \ell_q(\alpha)\ell_q(\tau) \leqslant \ell_q(\tau)\sqrt{\epsilon}$, hence $\ell_q(\tau) > \sqrt{\epsilon}$, so τ cannot belong to $C_q(\delta, \epsilon)$. $\square$

Given the quadratic differential q, let us denote as $N_q(\delta, T)$ the number of cylinders in the q-metric which have area bounded below by δ and length smaller than T. As Masur and Eskin showed, $N_q(\delta, T)$ grows quadratically as a function of T:

Theorem 3.2. *There exists $0 < \delta < 1$ and a constant $c_\delta > 0$ such that, for almost every quadratic differential q of unit area, we have*

$$\lim_{T\to\infty} \frac{N_q(\delta, T)}{T^2} = c_\delta$$

Proof. Let $0 < \delta < 1$. By the general counting argument of Eskin-Masur [EM01, Theorem 2.1] applied to the set of metric cylinders with area bounded below by δ, we get the existence of the limit c_δ almost everywhere. On the other hand, by [Mas93, Proposition 2.5], for every quadratic differential there exists some $\delta > 0$ such that $\liminf_{T\to\infty} \frac{N_q(\delta,T)}{T^2} > 0$, so the constant c_δ must be positive for some δ. $\square$

A finer statement, at least in the case of translation surfaces, is due to Vorobets [Vor05].

3.2 Asymptotic length of short curves

Let us now quantify the idea of keeping track of short curves in the flat metric. For the rest of the paper, we will fix some $\delta > 0$ for which Theorem 3.2 holds, and some $\epsilon < \delta$. Let us define the function $L : \mathcal{QM} \to \mathbb{R}$ as

$$L(q) := \sum_{\alpha\in C_q(\delta,\epsilon)} \frac{1}{\ell_q^2(\alpha)}$$

Note that by Lemma 3.1 the number of terms in the sum is always finite, so the function is well-defined. Let us fix denote by q_t the image of the quadratic differential q under the Teichmüller geodesic flow after time t. Our goal is to prove that the ergodic average of L is infinite:

Theorem 3.3. *For μ_{hol}-a.e. quadratic differential q of unit area, we have*

$$\lim_{T\to\infty} \frac{\int_0^T L(q_t)\ dt}{T} = \infty$$

In the proof of Theorem 3.3, we will make use of the following relations between metric cylinders and the geometry of Teichmüller discs. Let us fix a base point q_0 in the space of quadratic differentials, and call $\mathbb{D}$ the Teichmüller disc given by the $SL_2(\mathbb{R})$-orbit of q_0. For every metric cylinder α on q_0, there is an angle θ_α such that α is vertical in the quadratic differential $e^{i\theta_\alpha}q_0$. The angle θ_α determines a point in the circle at infinity of $\mathbb{D}$. For each metric cylinder on q_0 with core curve α, let us define the set

$$H_\epsilon(\alpha) := \{q \in \mathbb{D}\ :\ \ell_q^2(\alpha) \leqslant \epsilon\}$$

of points in the Teichmüller disc for which the length of α is less than the square root of ϵ. Recall the metric induced on $\mathbb{D}$ by the Teichmüller metric is the hyperbolic metric of constant curvature -4, and $H_\epsilon(\alpha)$ is a horoball for that metric.

Lemma 3.4. *The Euclidean diameter s of the horoball $H_\epsilon(\alpha)$ is*

$$s = \frac{2\epsilon}{\epsilon + \ell^2_{q_0}(\alpha)}$$

where $\ell_{q_0}(\alpha)$ is the length of α in the flat metric associated to the quadratic differential q_0.

Proof. By integrating the hyperbolic metric of curvature -4 we have

$$d(q_0, H_\epsilon(\alpha)) = \int_0^{1-s} \frac{dx}{1-x^2} = \frac{1}{2}\log\frac{2-s}{s}$$

and, since the Teichmüller map exponentially shrinks the curve α,

$$e^{-2d(q_0,H_\epsilon(\alpha))}\ell^2_{q_0}(\alpha) = \epsilon$$

hence the claim. ☐

We will need the following estimate from elementary Euclidean geometry:

Lemma 3.5. *In the unit disc, let $\theta(r,R)$ be the angle at the center of the disc corresponding to the intersection of the circle of radius $R \geqslant \frac{1}{2}$ centered at the origin, with a circle of radius $r \leqslant \frac{1}{2}$ tangent to the boundary, with $R + 2r - 1 \geqslant 0$. Then there is a constant K such that*

$$\frac{1}{K}\sqrt{(1-R)(R+2r-1)} \leqslant \theta(r,R) \leqslant K\sqrt{(1-R)(R+2r-1)}.$$

Proof. By the law of cosines, $r^2 = (1-r)^2 + R^2 - 2R(1-r)\cos(\theta/2)$. The claim follows by standard algebraic manipulation and approximation. ☐

Let $q_{t,\theta}$ denote the quadratic differential given by flowing the quadratic differential $e^{i\theta}q_0$ for time t.

Lemma 3.6. *For almost every quadratic differential q_0 there exists a constant $c > 0$, such that for each $\epsilon > 0$ there exists a time t_ϵ such that*

$$\sum_{\alpha \in C_{q_0}(\delta)} \mathrm{Leb}(\{\theta \in [0,2\pi] \; : \; q_{t,\theta} \in H_\epsilon(\alpha)\}) \geqslant c\epsilon \qquad \forall\, t \geqslant t_\epsilon,$$

where Leb *denotes Lebesgue measure on the circle.*

Proof. Let $\frac{1}{2} < R < 1$, and consider the set of horoballs of the collection $H_\epsilon(\alpha)$ with $\alpha \in C_{q_0}(\delta,\epsilon)$ and Euclidean diameter $s \geqslant \frac{3}{2}(1-R)$. By Lemma 3.4, these horoballs correspond precisely to metric cylinders with core curve α such that

$$\ell^2_{q_0}(\alpha) \leqslant \frac{3R+1}{3(1-R)}\epsilon$$

By Theorem 3.2, the number of such cylinders is, for R large, at least $\frac{c_\delta}{2}\frac{3R+1}{3(1-R)}\epsilon$. By Lemma 3.5, every corresponding horoball intersects the circle of Euclidean radius R centered at the origin in an arc of visual angle

$$\theta \geqslant \frac{1}{K\sqrt{2}}(1-R)$$

and by Lemma 3.1 every quadratic differential belongs to at most a universally bounded number M of horoballs, hence the total visual angle is at least $\frac{c_\delta}{6KM\sqrt{2}}\epsilon$. □

In order to prove Theorem 3.3, let us first define a discretized version of L. Namely, for each n and α we denote as $H_n(\alpha)$ the horoball

$$H_n(\alpha) := \{q \in \mathbb{D} \ : \ell_q^2(\alpha) \leqslant 2^{-n}\epsilon\}$$

Now, the function $\Psi : \mathcal{QM} \to \mathbb{R}$ is defined as

$$\Psi(q) := \sum_{\alpha \in C_q(\delta)} \sum_{n=1}^{\infty} 2^n \chi_{H_n(\alpha)}$$

It is easy to see that Ψ is bounded above by a multiple of L:

Lemma 3.7. *For each quadratic differential q, we have*

$$\Psi(q) \leqslant 4\epsilon L(q)$$

Proof. Let $\alpha \in C_q(\delta)$ be a short curve on q: then there exists a positive integer M such that

$$2^{-M}\epsilon \leqslant \ell_q^2(\alpha) \leqslant 2^{-M+1}\epsilon$$

Now, since q lies in $H_1(\alpha) \cup \cdots \cup H_M(\alpha)$,

$$\sum_{n=1}^{\infty} 2^n \chi_{H_n(\alpha)} \leqslant 1 + 2 + \cdots + 2^M \leqslant 2 \cdot 2^M \leqslant \frac{4\epsilon}{\ell_q^2(\alpha)}$$

and summing over α yields the claim. □

Proof of Theorem 3.3. By Lemma 3.7, it is enough to prove the statement for Ψ. Let us now truncate the function Ψ by defining, for each N,

$$\Psi_N(q) := \sum_{\alpha \in C_q(\delta)} \sum_{n=1}^{N} 2^n \chi_{H_n(\alpha)}$$

Let us now fix N. By Lemma 3.1, Ψ_N is bounded on the moduli space $\mathcal{MQ}$ of unit area quadratic differentials, hence μ_{hol}-integrable; by ergodicity of the geodesic flow, for a generic Teichmüller disc for almost all radial directions θ the ergodic average of Ψ_N along the flow tends to its integral:

$$\lim_{T\to\infty} \int_0^T \frac{\Psi_N(q_{t,\theta})\ dt}{T} = \int_{\mathcal{MQ}} \Psi_N(q)\ d\mu_{hol} \qquad \text{for a.e. } \theta$$

Then, if we integrate both sides w.r.t. to the angular measure $d\theta$ and apply the dominated convergence theorem,

$$\lim_{T\to\infty} \int_{S^1} d\theta \int_0^T \frac{\Psi_N(q_{t,\theta})\ dt}{T} = \int_{\mathcal{MQ}} \Psi_N(q)\ d\mu_{hol}$$

and by Fubini

$$\lim_{T\to\infty} \frac{\int_0^T dt \int_{S^1} \Psi_N(q_{t,\theta})\ d\theta}{T} = \int_{\mathcal{MQ}} \Psi_N(q)\ d\mu_{hol}$$

Now, by Lemma 3.6, for each $t > T_{2^{-N}}$

$$\int_{S^1} \Psi_N(q_{t,\theta})\ d\theta \geqslant \sum_{n=1}^{N} 2^n \cdot c2^{-n} = cN$$

hence

$$\int_{\mathcal{MQ}} \Psi_N(q)\ d\mu_{hol} = \lim_{T\to\infty} \frac{\int_0^T dt \int_{S^1} \Psi_N(q_{t,\theta})\ d\theta}{T} \geqslant cN$$

Since the previous estimate works for all N, then also

$$\int_{\mathcal{MQ}} L(q)\ d\mu_{hol} = \infty$$

hence the ergodic average tends to infinity almost everywhere:

$$\lim_{T\to\infty} \int_0^T \frac{L(q_t)\ dt}{T} = \int_{\mathcal{MQ}} L(q)\ d\mu_{hol} = \infty \qquad \text{for a.e. } q \in \mathcal{MQ}.$$

□

3.3 Average excursion

Let us now turn the asymptotic estimate of the previous section into an asymptotic about excursions. If q is a quadratic differential, let us denote as γ_q the corresponding Teichmüller geodesic ray. We now define the concept of *total excursion* traveled by the geodesic γ_q inside the horoballs up to time T:

Definition 3.8. *Given a quadratic differential q, the* total excursion *$E(q,T)$ is the sum of all excursions in all horoballs crossed by the geodesic ray γ_q up to time T:*

$$E(q,T) := \sum_{\gamma_q([0,T])\cap H_\epsilon(\alpha)\neq\emptyset} E(\gamma_q, H_\epsilon(\alpha))$$

Our goal is to prove that also the average total excursion is infinite.

Theorem 3.9. *For μ_{hol}-almost every quadratic differential q of unit area,*

$$\lim_{T\to\infty} \frac{E(q,T)}{T} = \infty$$

Theorem 3.9 follows from Theorem 3.3 and the following

Proposition 3.10. *Let q be a quadratic differential with geodesic ray γ_q, and let $T > 0$ be such that both q and $\gamma_q(T)$ lie outside all horoballs of the type $H_\epsilon(\alpha)$. Then*

$$\int_0^T L(q_t)\ dt \leqslant \frac{C}{\epsilon} E(q,T)$$

for some universal constant C.

Proof. Let $\alpha \in C_q(\delta)$ be a curve which has become short before time T, i.e. such that $\gamma_q([0,T]) \cap H_\epsilon(\alpha)$ is non-empty. Let T_1 be the time the geodesic enters $H_\epsilon(\alpha)$, and T_2 the time the geodesic exits. Moreover, let N be the maximum integer k such that the geodesic enters $H_k(\alpha)$. Note that there is a universal constant C_1 such that for each $n \geqslant 1$ and each α

$$\text{Leb}(\{t \in [0,T]\ :\ q_t \in H_{n+1}(\alpha) \setminus H_n(\alpha)\}) \leqslant C_1$$

Then

$$\int_{T_1}^{T_2} \frac{1}{\ell^2_{q_t}(\alpha)}\,dt \leqslant \sum_{n=1}^{N} \frac{2^n}{\epsilon}\mathrm{Leb}(\{t \in [0,T] \,:\, q_t \in H_{n+1}(\alpha)\backslash H_n(\alpha)\}) \leqslant \frac{C_1 \cdot 2^{N+1}}{\epsilon}$$

In order to compare the right hand side with the excursion, let us denote by $\tilde{\epsilon}$ the smallest value of $\ell^2_{q_t}(\alpha)$ along the geodesic ray γ_q. By the definition of N, we have $\tilde{\epsilon} \asymp 2^{-N}\epsilon$. Now, by the definition of excursion and Lemma 2.8,

$$E(\gamma_q, H_\epsilon(\alpha)) = \frac{\phi_{max}}{\phi_0} \asymp \frac{\sin\phi_{max}}{\sin\phi_0} = \frac{\epsilon}{\tilde{\epsilon}} \asymp 2^N$$

(where all the approximate equalities hold up to multiplicative constants), hence the claim follows. ☐

3.4 The word metric

Let us complete the proof of Theorem 1.3 by proving that the word metric is bounded below by the total excursion. Let us pick ϵ_0 to define the thick part as in section 2.5, and let us choose δ so that Theorem 3.2 holds. Finally, we choose ϵ so that if X belongs to the thick part $\mathcal{T} \setminus \mathcal{T}_{\epsilon_0}$ and α is the core curve of a metric cylinder of q-area larger than δ on X, then $\ell^2_q(\alpha) \geqslant \epsilon$.

Let X_0 lie in the thick part $\mathcal{T}\setminus\mathcal{T}_{\epsilon_0}$, and let γ be a Teichmüller geodesic with $\gamma(0) = X_0$. Recall for each time t, h_t is a closest point projection of $\gamma(t)$ to the Teichmüller lattice.

Proposition 3.11. *If $\gamma(T)$ lies in the thick part $\mathcal{T} \setminus \mathcal{T}_{\epsilon_0}$, then*

$$d_G(1, h_T) \geqslant C_1 E(\gamma, T) - C_2$$

where the constants depend only on X_0, the choice of ϵ_0 and the choice of generating set for $Mod(S)$.

Proof. Since $\gamma(0)$ and $\gamma(T)$ lie in the thick part, by Lemma 2.13

$$d_G(1, h_T) \asymp d_M(m_0, m_T)$$

By the Masur-Minsky quasi-distance formula (Theorem 2.2), for any B large enough

$$d_M(m_0, m_T) \asymp \sum_{Y \subseteq S} \lfloor d_Y(m_0, m_T) \rfloor_B \geqslant \sum_{\gamma([0,T]) \cap H_\epsilon(\alpha) \neq \emptyset} \lfloor d_\alpha(m_0, m_T) \rfloor_B$$

where on the right-hand side we only consider projections to annuli of area bounded below, and whose core curve becomes short before time T. Now by Proposition 2.9, for some constants K_1 and K_2,

$$d_\alpha(m_0, m_T) \geqslant K_1 E(\gamma, H_\epsilon(\alpha)) - K_2$$

so if $B \geqslant K_2$

$$\lfloor d_\alpha(m_0, m_T) \rfloor_B \geqslant \lfloor K_1 E(\gamma, H_\epsilon(\alpha)) - K_2 \rfloor_B \geqslant \frac{K_1}{2} \lfloor E(\gamma, H_\epsilon(\alpha)) \rfloor_{\frac{2B}{K_1}}$$

and we can choose $\tilde{\epsilon}$ a bit smaller than ϵ so that $\lfloor E(\gamma, H_\epsilon(\alpha) \rfloor_{\frac{2B}{K_1}} \geqslant E(\gamma, H_{\tilde{\epsilon}}(\alpha))$, hence

$$\sum_{\gamma([0,T]) \cap H_\epsilon(\alpha) \neq \emptyset} \lfloor d_\alpha(m_0, m_T) \rfloor_B \geqslant \sum_{\gamma([0,T]) \cap H_{\tilde{\epsilon}}(\alpha) \neq \emptyset} E(\gamma, H_{\tilde{\epsilon}}(\alpha)) = E(q, T)$$

☐

Proof of Theorem 1.3. By Theorem 2.3, the relative metric is a lower bound for Teichmüller distance, i.e. there exist constants K, c, depending only on the topology of S, such that

$$d_{rel}(1, h_T) \leqslant KT + c$$

Theorem 1.3 then follows from Theorem 3.9 and Proposition 3.11. $\square$

4 Hitting measure sampling

Let μ be a finitely-supported measure on the mapping class group $G = \mathrm{Mod}(S)$. The *step space* is the infinite product $G^{\mathbb{N}}$ with the product measure $\mathbb{P} := \mu^{\mathbb{N}}$. Let $w_n = g_1 g_2 \ldots g_2$ be the location of the random walk after n steps. The *path space* is $G^{\mathbb{N}}$, with the pushforward of the product measure under the map

$$(g_1, g_2, g_3, \ldots) \mapsto (w_1, w_2, w_3, \ldots).$$

It will also be convenient to consider *bi-infinite* sample paths. In this case the step space is the set $G^{\mathbb{Z}}$ of bi-infinite sequences of group elements with the product measure. The location of the random walk is given by $w_0 = 1$, and $w_n = g_1 g_2 \ldots g_n$ if n, if n is positive, and $w_n = g_0^{-1} g_{-1}^{-1} \ldots g_{n-1}^{-1}$, if n is negative. The path space is $G^{\mathbb{Z}}$, as a set, but with measure coming from the pushforward of $\mathbb{P}$ under the map

$$(\ldots, g_{-1}, g_0, g_1, g_2, \ldots) \mapsto (\ldots w_{-1}, w_0, w_1, w_2, \ldots).$$

Let us fix a base point $X_0 \in \mathcal{T}$, and consider the image of the sample paths $w_n X_0$ in $\mathcal{T}$. Kaimanovich and Masur showed that almost every sample path converges to a uniquely ergodic foliation in the space $\mathcal{PMF}$ of projective measured foliations, Thurston's boundary for Teichmüller space. Recall that the *harmonic measure* ν on $\mathcal{PMF}$ is defined as the hitting measure of the random walk, i.e. for any measurable subset $A \subseteq \mathcal{PMF}$,

$$\nu(A) := \mathbb{P}(w_n \ : \lim_{n \to \infty} w_n X_0 \in A)$$

Theorem 4.1 (Kaimanovich and Masur [KM96])**.** *Let μ be a probability distribution on the mapping class group whose support generates a non-elementary subgroup. Then almost every sample path $(w_n)_{n \in \mathbb{N}}$ converges to a uniquely ergodic foliation in $\mathcal{PMF}$, and the resulting hitting measure ν is the unique non-atomic μ-stationary measure on $\mathcal{PMF}$.*

The mapping class group is finitely generated, and let d_G be the word metric on the mapping class group with respect to some choice of generating set. Since the mapping class group is non-amenable, the random walk makes linear progress in the word metric d_G, or indeed in any metric quasi-isometric to the word metric.

Theorem 4.2 (Kesten [Kes59], Day [Day64])**.** *Let μ be a probability distribution on a group, whose support generates a non-amenable subgroup. Then there exists a constant $c_1 > 0$ such that for almost all sample paths*

$$\lim_{n \to \infty} \frac{d_G(1, w_n)}{n} = c_1. \tag{13}$$

Even though the relative metric is smaller than the word metric, a more recent result proves that the growth rate is still linear in the number of steps.

Theorem 4.3 ([Mah10a]). *Let μ be a probability distribution on the mapping class group, whose support generates a non-elementary subgroup. Then there is a constant $c_2 > 0$ such that for almost all sample paths*

$$\lim_{n\to\infty} \frac{d_{rel}(1, w_n)}{n} = c_2$$

From these results it follows that the quotient between the word metric and the relative metric converges to c_1/c_2 for almost every sample path, i.e.

$$\lim_{n\to\infty} \frac{d_G(1, w_n)}{d_{rel}(1, w_n)} = \frac{c_1}{c_2}$$

for almost all sample paths. The limit above is a limit taken along the locations $(w_n)_{n\in\mathbb{N}}$ of the random walk. In order to compare this to the previous statistic we need to relate locations of the random walk to points on a Teichmüller geodesic.

Because of Kaimanovich-Masur, for almost every bi-infinite sample path $w \in G^{\mathbb{Z}}$, there are well-defined maps

$$F^{\pm} : G^{\mathbb{Z}} \to \mathcal{PMF}$$

given by

$$F^{+}(w) := \lim_{n\to\infty} w_n X_0$$

and

$$F^{-}(w) := \lim_{n\to\infty} w_{-n} X_0.$$

Furthermore, the two foliations $F^{+}(w)$ and $F^{-}(w)$ are almost surely uniquely ergodic and distinct, so there is a unique oriented Teichmüller geodesic γ_w whose forward limit point in $\mathcal{PMF}$ is $F^{+}(w)$ and whose backward limit point is $F^{-}(w)$. There is also a unique geodesic ray ρ_w starting at the basepoint X_0 whose forward limit point is F^{+}. We shall always parameterize ρ_w as unit speed geodesic with $\rho_w(0) = X_0$. As F^{+} is uniquely ergodic, the distance between γ_w and ρ_w tends to zero [Mas80], and we shall parameterize γ_w such that $d_{\mathcal{T}}(\rho_w(t), \gamma_w(t)) \to 0$.

For each bi-infinite sample path we can define the function

$$D : G^{\mathbb{Z}} \to \mathbb{R}$$

given by

$$D(w) := d_{\mathcal{T}}(X_0, \gamma_w)$$

which represents the Teichmüller distance between the base point X_0 and the geodesic γ_w. This is well-defined and measurable, by Lemma 1.4.4 of [KM96]. In particular, this implies that for any $\epsilon > 0$ there is a constant M such that the probability that $D(w) \leqslant M$ is at least $1 - \epsilon$.

The shift map σ maps the step space to itself by incrementing the index of each step by one, i.e.

$$\sigma\colon (g_n)_{n\in\mathbb{Z}} \mapsto (g_{n+1})_{n\in\mathbb{Z}}.$$

This is a measure preserving ergodic transformation on the step space, and the induced action of σ on the path space is given by

$$\sigma\colon (w_n)_{n\in\mathbb{Z}} \mapsto (w_1^{-1} w_{n+1})_{n\in\mathbb{Z}}.$$

4.1 Distance between geodesic and sample path

The geodesic γ_w is determined by its endpoints $F^+(w)$ and $F^-(w)$, and the distribution of these pairs is given by harmonic measure ν and reflected harmonic measure $\check{\nu}$ respectively.

The distance from a location w_n to the corresponding geodesic γ_w is given by

$$d_{\mathcal{T}}(w_n X_0, \gamma_w) = d_{\mathcal{T}}(X_0, w_n^{-1}\gamma_w)$$

since the mapping class group acts on $\mathcal{T}$ by isometries, and by the definition of the shift map,

$$d_{\mathcal{T}}(w_n X_0, \gamma_w) = d_{\mathcal{T}}(X_0, \gamma_{\sigma^n w}).$$

As already noted in [KM96], if ϵ is sufficiently small, almost every geodesic with respect to harmonic measure returns to the ϵ-thick part $\mathcal{T} \setminus \mathcal{T}_\epsilon$ infinitely often.

Our goal is to show that every step of the random walk lies within sublinear distance in the word metric from some point in the thick part of the limit geodesic. In [Tio12], sublinear tracking is proven in the Teichmüller metric: we will adapt the argument to the word metric. First, we give a general criterion which guarantees that a sequence of real numbers $(d_n)_{n\in\mathbb{N}}$ grow sublinearly. We then apply this to our set up, where the sequence d_n will correspond to distance between locations of the random walk and certain lattice points close to the corresponding Teichmüller geodesic ray, in either the word metric, or the Teichmüller metric.

Lemma 4.4 ([Tio12])**.** *Let $d_n \geqslant 0$ be a sequence of non-negative real numbers, such that*

1. $|d_n - d_{n+1}| \leqslant \ell$ *for some* $\ell > 0$
2. *For each* $p > 0$*, there exists* $M \geqslant 0$ *such that*

$$\limsup_{n\to\infty} \frac{\#\{k : 1 \leqslant k \leqslant n, d_k \geqslant M\}}{n} \leqslant p$$

Then

$$\lim_{n\to\infty} \frac{d_n}{n} = 0$$

Proof. Suppose on the contrary that there exists $c > 0$ and a subsequence $\{n_k\}$ such that $d_{n_k} \geqslant c n_k$. Let now M be any real number. By (1)

$$d_{n_k - h} \geqslant d_{n_k} - h\ell \geqslant c n_k - h\ell$$

hence by choosing $n_k \geqslant \frac{2M}{c}$ we have

$$d_{n_k - h} \geqslant \frac{c}{2} n_k \geqslant M \qquad \text{for each } 0 \leqslant h \leqslant \frac{c n_k}{2\ell}$$

hence there exists a set $A \subseteq \mathbb{N}$ of density at least $\frac{c}{2\ell}$ such that $d_k \geqslant M$ for $k \in A$. By choosing some $p < \frac{c}{2\ell}$ we contradict condition (2). $\square$

We now explain how to apply the lemma above in the current setting. Given a point $X \in \mathcal{T}$, let us denote as $\operatorname{proj}(X)$ the set of lattice points at minimal distance from X:

$$\operatorname{proj}(X) := \{h \in G : d_{\mathcal{T}}(hX_0, X) \text{ is minimal}\}$$

Such a projection may possibly vary wildly if X lies in the thin part, but it is controlled in the thick part: namely, given $\epsilon > 0$ there is a constant $K(\epsilon)$ such that

$$d_{\mathcal{T}}(X, hX_0) \leqslant K(\epsilon), \qquad \forall X \notin \mathcal{T}_\epsilon \quad \forall h \in \mathrm{proj}(X)$$

We now associate to almost every sample path w a subset $P(w)$ of the mapping class group, which we now describe. Almost every bi-infinite sample path $w \in G^{\mathbb{Z}}$ determines two uniquely ergodic foliations, $F^{\pm}(w)$. Let γ_w be the bi-infinite Teichmüller geodesic joining them. Now, let us define $P(w)$ as the set of mapping class group elements $h \in G$ such that hX_0 is the closest projection from some point X in $\gamma_w \setminus \mathcal{T}_\epsilon$, i.e.

$$P(w) := \bigcup_{X \in \gamma_w \setminus \mathcal{T}_\epsilon} \mathrm{proj}(X).$$

This is illustrated in Figure 5.



Figure 5: Sample path locations and basepoint orbits close to the geodesic.

The key result is the following:

Proposition 4.5. *Fix $\epsilon > 0$, sufficiently small. Then for almost every sample path $(w_n)_{n\in\mathbb{N}}$, with corresponding Teichmüller ray ρ_w, there exists a sequence of times $t_n \to \infty$ with $\rho_w(t_n) \in \rho_w \setminus \mathcal{T}_\epsilon$, such that*

$$\lim_{n\to\infty} \frac{d_G(w_n, h_n)}{n} = 0$$

for any $h_n \in \mathrm{proj}(\rho_w(t_n))$.

Proof. Let us fix $\epsilon > 0$ sufficiently small. Recall that $P(w)$ is the collection of group elements corresponding to closest lattice points to points on the geodesic γ_w which lie in the thick part of Teichmüller space. Note that, since the mapping class group acts by isometries with respect to both the Teichmüller and word metrics, then P is equivariant, in the sense that

$$P(\sigma^n w) = w_n^{-1} P(w)$$

Let us now define the function $\varphi : G^{\mathbb{Z}} \to \mathbb{R}$ on the space of bi-infinite sample paths as

$$\varphi(w) := d_G(1, P(w))$$

i.e. the minimal word-metric distance between the base point X_0 and the set of closest projections from the thick part of the geodesic γ_w. The shift map $\sigma : G^{\mathbb{Z}} \to G^{\mathbb{Z}}$ acts on the space of sequences, ergodically with respect to the product measure $\mu^{\mathbb{Z}}$. By the equivariance of P, we have

$$\varphi(\sigma^n w) = d_G(w_n, P(w))$$

Since φ attains finite values almost everywhere, for each $p > 0$ one can choose $M > 0$ such that

$$\mathbb{P}(w \in G^{\mathbb{Z}} : \varphi(w) \geqslant M) \leqslant p$$

Hence, by ergodicity, for almost all sample paths w,

$$\lim_{n\to\infty} \frac{\#\{1 \leqslant k \leqslant n : \varphi(\sigma^k w) \geqslant M\}}{n} \leqslant p$$

Now, since the measure is finitely-supported, then the function φ has bounded increments; indeed,

$$|\varphi(\sigma^n w) - \varphi(\sigma^{n+1} w)| = |d_G(1, P(\sigma^n w)) - d_G(1, P(\sigma^{n+1} w))|,$$

and by the equivariance of P, this is equal to

$$|d_G(w_n, P(w)) - d_G(w_{n+1}, P(w))| \leqslant d_G(w_n, w_{n+1}) \leqslant D,$$

where D is the diameter of the support of the measure μ. Then by Lemma 4.4, setting $d_n := \varphi(\sigma^n w)$, it follows that for almost all bi-infinite paths w

$$\lim_{n\to\infty} \frac{d_G(w_n, P(w))}{n} = 0$$

By definition of $P(w)$, there exists a sequence of times t_n, such that $\gamma_w(t_n)$ lies in $\gamma_w \setminus \mathcal{T}_\epsilon$, the ϵ-thick part of the geodesic γ_w, and group elements $p_n \in G$ such that $p_n \in \mathrm{proj}(\gamma_w(t_n))$, and furthermore

$$\lim_{n\to\infty} \frac{d_G(w_n, p_n)}{n} = 0. \tag{14}$$

Now let F^+ be the terminal foliation of the geodesic γ_w, and denote as ρ_w the geodesic ray through X_0 with terminal foliation F^+. We have obtained a sequence of points lying in the intersection of the geodesic γ_w with the thick part $\mathcal{T} \setminus \mathcal{T}_\epsilon$, and we now show how to obtain a sequence of points lying in the intersection of the geodesic ρ_w with the thick part $\mathcal{T} \setminus \mathcal{T}_\epsilon$.

Recall that since γ_w and ρ_w have the same terminal foliation F^+, and F^+ is almost surely uniquely ergodic, then the distance between the positive ray ρ_w and the geodesic γ_w tends to zero, and we have chosen parameterizations such that $d_{\mathcal{T}}(\gamma_w(t), \rho_w(t)) \to 0$. In particular, after discarding finitely many initial values, we may assume

$$d_{\mathcal{T}}(\gamma_w(t_n), \rho_w(t_n)) \leqslant \frac{\log 2}{2},$$

for all n. Now for each n sufficiently large consider the sequence take $\rho_w(t_n)$. Then:

1. By Wolpert's lemma, $\rho_w(t_n)$ lies in the $\frac{\epsilon}{2}$-thick part;
2. if $h_n \in \mathrm{proj}(\rho_w(t_n))$, then $d_{\mathcal{T}}(\rho_w(t_n), h_n X_0) \leqslant K(\frac{\epsilon}{2})$ so

   $$\begin{aligned} d_{\mathcal{T}}(h_n X_0, p_n X_0) &\leqslant d_{\mathcal{T}}(h_n X_0, \rho_w(t_n)) + d_{\mathcal{T}}(\rho_w(t_n), \gamma_w(t_n)) + d_{\mathcal{T}}(\gamma_w(t_n), p_n X_0) \\ &\leqslant 1 + 2K(\frac{\epsilon}{2}), \end{aligned}$$

   hence $d_G(h_n, p_n) \leqslant K'$, so by eq. (14) we have also $\lim d_G(w_n, h_n)/n \to 0$.

This completes the proof of Proposition 4.5. ☐

4.2 Intermediate times

So far, we have shown that every step of the sample path is close enough to some point on the thick part of the geodesic, hence the closest projection to the lattice will behave like the sample path. However, we still need to deal with the case in which there are points in the thick part of the Teichmüller geodesic which are not close to the sample path.



Figure 6: Intermediate times.

Proof of Theorem 1.3. Given a sample path w, let ρ_w be the geodesic ray joining the base point X_0 to the limit foliation $F^+(w)$, and let t_n be the sequence of times given by Proposition 4.5. Let now $T > 0$ be a time for which the geodesic $\rho_w(T)$ lies in the thick part, and let $h_T X_0$ be a projection of $\rho_w(T)$ to the Teichmüller lattice. Since $t_n \to \infty$, there exists an index $n = n(T)$ such that $t_n \leqslant T \leqslant t_{n+1}$. By Proposition 2.14, there exist constants $C_1 > 0$, C_2 such that

$$d_G(h_n, h_T) \leqslant C_1 d_G(h_n, h_{n+1}) + C_2$$

Moreover, by Proposition 4.5 and triangle inequality,

$$\lim_{n\to\infty} \frac{d_G(h_n, h_{n+1})}{n} \leqslant \lim_{n\to\infty} \frac{d_G(h_n, w_n) + d_G(w_n, w_{n+1}) + d_G(w_{n+1}, h_{n+1})}{n} = 0$$

Thus, we also have

$$\lim_{n\to\infty} \frac{d_G(h_n, h_T)}{n} = 0$$

and again by Proposition 4.5

$$\lim_{n\to\infty} \frac{d_G(w_n, h_T)}{n} \leqslant \lim_{n\to\infty} \frac{d_G(w_n, h_n) + d_G(h_n, h_T)}{n} = 0$$

Similarly, since the relative metric is bounded above by the word metric,

$$\lim_{n\to\infty} \frac{d_{rel}(w_n, h_T)}{n} = 0$$

Finally, by computing the ratio between the word and relative metric,

$$\lim_{\substack{T\to\infty \\ \rho_w(T)\notin\mathcal{T}_\epsilon}} \frac{d_G(1, h_T)}{d_{rel}(1, h_T)} = \lim_{\substack{T\to\infty \\ \rho_w(T)\notin\mathcal{T}_\epsilon}} \frac{\frac{d_G(1,h_T)}{n(T)}}{\frac{d_{rel}(1,h_T)}{n(T)}} = \lim_{n\to\infty} \frac{\frac{d_G(1,w_n)}{n}}{\frac{d_{rel}(1,w_n)}{n}} = \frac{c_1}{c_2} > 0$$

This completes the proof of Theorem 1.3. ☐

5 Fuchsian groups

Let G be a lattice in $PSL(2,\mathbb{R})$ such that the quotient $X = G\backslash\mathbb{H}^2$ is a finite area non-compact orbifold. Fix a thick-thin decomposition of X such that the thin parts correspond to disjoint cusps $c_1, \cdots, c_p$ of X, and let $g_1, \ldots, g_p$ be the generators of the corresponding parabolic subgroups G_i of G.

A choice of generators $\mathcal{A}$ for G defines a *word metric* on G and also a *relative metric* on G which is the word metric with respect to the larger set of generators $\mathcal{A}' = \mathcal{A} \cup \{g_i^n\}$ i.e., along with the generators of G, the set $\mathcal{A}'$ includes all powers of all the parabolic generators g_i. It is well known that G with the relative metric d_{rel} is Gromov hyperbolic. In fact, G is strongly hyperbolic relative to the parabolic subgroups G_i ([Far98, Theorem 4.11]).

Let μ be a finitely supported measure on G, such that the support of μ generates a non-elementary subgroup of G i.e., the subgroup contains a pair of hyperbolic isometries with different fixed points. The random walk on G defined by μ is the G-valued stochastic process $w_1, w_2, \ldots$ with

$$\begin{cases} w_1 := g_1 \\ w_{n+1} := w_n g_{n+1} \end{cases}$$

where each g_i is drawn from G with probability μ and independently of the previous ones.

Given a basepoint $x_0 \in \mathbb{H}^2$, one can consider the projection in $\mathbb{H}^2$ of the random walk by the action of G, i.e. the sequence $(w_1(x_0), w_2(x_0), \ldots, w_n(x_0), \ldots)$. Then a theorem of Furstenberg [Fur63] assures that for almost all sequences $w = (g_1, g_2, \ldots)$ (with respect to the product measure $\mu^{\mathbb{N}}$) the random walk converges to some point $p_\infty \in S^1 = \partial\mathbb{H}^2$. The harmonic measure ν on the boundary records the probability that the random walk hits a particular part of $\partial\mathbb{H}^2$, i.e.,

$$\nu(A) = \text{Prob}\left(\lim_{n\to\infty} w_n(p_0) \in A\right).$$

We now provide a quick outline of the proof of Theorem 1.1. Since the proof in this case is much simpler than for mapping class groups we leave it to the reader to reconstruct the exact argument.

The unit tangent bundle $T^1\mathbb{H}^2$ carries the *Liouville measure* which in the upper half-plane model is given by $d\ell = \frac{dx\ dy\ d\theta}{y^2}$ and is invariant by the action of $PSL_2(\mathbb{R}) = \mathrm{Isom}(\mathbb{H}^2)$. Hence it descends to a geodesic flow invariant measure on the unit tangent bundle to $X = G\backslash\mathbb{H}^2$. Moreover, it is classically known (Hopf) that such flow is ergodic (and indeed mixing). The conditional measure on the unit ball in the tangent space at any point is the pullback via the visual map of the standard Lebesgue measure on $\partial\mathbb{H}^2 = S^1$.

The cusps $c_1, \ldots, c_p$ lift to $\mathbb{H}^2$ to give a horoball packing $\mathcal{H}$. By studying this packing, Sullivan [Sul82] proved that a geodesic ray generic with respect to the Lebesgue measure is recurrent to the thick part of X venturing into the cusps infinitely often with maximum depth in the cusps of about $\log t$ where t is the time along the geodesic ray. Sullivan's theorem is a precursor of Masur's approach in Teichmüller space establishing that every Teichmüller disk has a horoball packing with similar good properties.

Given a horoball $H \in \mathbb{H}^2$ and a geodesic γ that enters and leaves H, we define as earlier, the excursion $E(\gamma, H)$ to be the distance in the path metric on ∂H between the entry and exit points. Sullivan's theorem implies that a lift in $\mathbb{H}^2$ of a Lebesgue typical geodesic ray enters and leaves infinitely many horoballs in the packing. We use this setup to estimate from below the word length along a Lebesgue typical geodesic in terms of the sum of the excursions in these horoballs.

For a time t along such a geodesic ray γ one can keep track of which image under G of a fixed fundamental domain does the point γ_t lie in. Up to an element of the finite stabilizer in G of the fundamental domain, this associates to γ_t a group element h_t. Coarsely, h_t is the group element given by a point in the orbit Gx_0 that is the closest to γ_t.

Proposition 5.1. *There exists uniform constants $c, d > 0$ such that*

$$d_G(1, h_t) \geqslant c \left(\sum_{\gamma([0,t]) \cap H \neq \emptyset} E(\gamma, H) \right) - d$$

Proof. A finite index subgroup of G is quasi-isometric to G. Passing to a finite index subgroup, we can assume that G has no torsion.

The group G acts properly discontinuously and cocompactly on the thick part $\mathbb{H}^2 \setminus \cup_{\mathcal{H}} H$. By Milnor-Svarc lemma, the Cayley graph of G is quasi-isometric to the thick part with the path metric. There is a unique geodesic word representing h_t and in this case it is easy to verify that the sum of the excursions till t is a coarse lower bound on $d_G(1, h_t)$ (See [Far98] Lemma 4.5, 4.8 and 4.9). Combining all the quasi-isometry constants yields the proposition. ☐

Inside $H \in \mathcal{H}$, let H_n be the horoball consisting of points that are at least $\log n$ deep in H in the hyperbolic metric:

$$H_n := \{x \in \mathbb{H}^2 \ : \ d(x, \partial H) \geqslant \log n\}$$

Let us denote as X_n the quotient of H_n under the action of G, that is a subset of X. By the ergodicity of geodesic flow, the proportion of time that a geodesic ray spends in X_n is asymptotically the same as the volume of $T^1 X_n$, and this volume can be easily calculated to be $1/n$ up to a uniform multiplicative constant. Let χ_n be the characteristic function of $T^1(X_{2^n})$, and let $\psi : T^1\mathbb{H}^2 \to \mathbb{R}$ be

$$\psi(v) := \sum_{n=1}^{\infty} 2^n \chi_n(v)$$

The function is well defined since for any v there is some N such that $v \in T^1X_{2^N} \setminus T^1X_{2^{N+1}}$. As a consequence of the $1/n$ decay of volumes, the ergodic average of ψ is infinite:

Proposition 5.2. *For almost every tangent vector $v \in T^1\mathbb{H}^2$ with respect to Liouville measure,*

$$\lim_{T\to\infty} \int_0^T \frac{\psi(v(\gamma_t))}{T} dt = \infty \tag{15}$$

where $v(\gamma_t)$ is the unit tangent vector to the geodesic γ originating from v at the point γ_t.

Proof. Let ψ_N be the truncation:

$$\psi_N(v) = \sum_{n=1}^{N} 2^n \chi_n(v)$$

Up to a uniform multiplicative constant,

$$\int_{T^1X} \psi_N \; d\ell \asymp N$$

By ergodicity, along ℓ-almost every geodesic ray γ

$$\lim_{T\to\infty} \int_0^T \frac{\psi_N(v(\gamma_t))}{T} dt = \int_{T^1X} \psi_N \; d\ell \asymp N$$

where $v(\gamma_t)$ is the unit tangent vector to γ at the point γ_t. As a consequence, along ℓ-almost every geodesic ray γ the inequality

$$\lim_{T\to\infty} \int_0^T \frac{\psi(v(\gamma_t))}{T} dt \geqslant \lim_{T\to\infty} \int_0^T \frac{\psi_N(v(\gamma_t))}{T} dt \asymp N$$

holds for all N, which yields the claim. ☐

Proposition 5.3. *Let H be a horoball and let $t_1 < t_2$ be the entry and exit times in H for a geodesic ray γ. Then up to uniform additive and multiplicative constants*

$$\int_{t_1}^{t_2} \psi(v(\gamma_t)) \; dt \asymp E(\gamma, H)$$

Proof. Let N be the smallest number such that $\psi(v(\gamma_t)) = \psi_N(v(\gamma_t))$. Then up to a uniform additive constant $2^N \leqslant E(\gamma, H) \leqslant 2^{N+1}$. For $k \leqslant N-1$, there are uniform lower and upper bounds independent of k and N for the amount of time s_k that a geodesic ray γ can spend in $H_{2^k} \setminus H_{2^{k+1}}$. There is also a uniform upper bound independent of N for the amount of time s_N that the ray γ can spend in $H_{2^N} \setminus H_{2^{N+1}}$. These bounds imply

$$\int_{t_1}^{t_2} \psi_N(v(\gamma_t)) \; dt \asymp \sum_{k=1}^{N} s_k \left(\sum_{j=1}^{k} 2^j \right) \asymp 2^N \asymp E(\gamma, H)$$

☐

Combining Propositions 5.1, 5.3 and Equation (15) we have

$$\lim_{t\to\infty}\frac{d_G(1,h_t)}{t} = +\infty$$

On the other hand, the relative length of h_t is up to a uniform multiplicative constant bounded above by t. In fact, by ergodicity, the ray γ spends a definite proportion of its time in the thick part of X. This implies that the relative length of h_t grows linearly in t. Combining this observation with the limit above proves the first part of Theorem 1.2.

For the harmonic measure, we leverage the following facts. Since G is non-amenable, a random walk makes linear progress in the word metric and also in the hyperbolic metric for the action of G on $\mathbb{H}^2$. Since G with the relative metric is δ-hyperbolic, the random walk also makes linear progress in the relative metric. This can be proved mimicking the argument in [Mah10a]. Hence by taking the quotient, the limit

$$\lim_{n\to\infty}\frac{d_G(1,w_n)}{d_{rel}(1,w_n)}$$

exists (and is finite) along almost every sample path $w = (w_1, w_2, \dots)$.

Since almost every bi-infinite sample path converges to a pair of distinct points (λ_1,λ_2) in S^1 one can consider the geodesic in $\mathbb{H}^2$ joining λ_1 and λ_2 and consider its distance from the basepoint. This defines a measurable function on the space of bi-infinite sample paths. So there exists $D > 0$ such that the set of bi-infinite sample paths for which the distance of the geodesic from the basepoint is less than D has positive measure. Ergodicity of the shift map then implies that almost every sample path gets within distance D of the geodesic ray it tracks infinitely often. Moreover, it is well known that in the case of Fuchsian groups sample paths track geodesics sub-linearly. Let w_{n_i} be the subsequence that is within distance D of γ and let γ_{t_i} be the corresponding closest points on γ. The limit along t_i of $d_G(1,h_t)/d_{rel}(1,h_t)$ exists. Sub-linear tracking implies that the limit $d_G(1,h_t)/d_{rel}(1,h_t)$ equals the limit along the subsequence t_n and this proves the second part of Theorem 1.2.

References

[Beh06] Jason A. Behrstock, *Asymptotic geometry of the mapping class group and Teichmüller space*, Geom. Topol. **10** (2006), 1523–1578.

[Day64] Mahlon Marsh Day, *Convolutions, means, and spectra*, Illinois J. Math. **8** (1964), 100–111.

[DDM11] Spencer Dowdall, Moon Duchin, and Howard A. Masur, *Statistical hyperbolicity in Teichmüller space* (2011), available at `arXiv:1108.5416`.

[EM01] Alex Eskin and Howard A. Masur, *Asymptotic formulas on flat surfaces*, Ergodic Theory Dynam. Systems **21** (2001), no. 2, 443–478.

[Far98] Benson Farb, *Relatively hyperbolic groups*, Geom. Funct. Anal. **8** (1998), no. 5, 810–840.

[Fur63] Harry Furstenberg, *Noncommuting random products*, Trans. Amer. Math. Soc. **108** (1963), 377–428.

[Gad09] Vaibhav Gadre, *Harmonic measures for distributions with finite support on the mapping class group are singular* (2009), available at `arXiv:0911.2891`.

[GLJ90] Yves Guivarc'h and Yves Le Jan, *Sur l'enroulement du flot géodésique*, C. R. Acad. Sci. Paris Sér. I Math. **311** (1990), no. 10, 645–648.

[GLJ93] Yves Guivarc'h and Yves Le Jan, *Asymptotic winding of the geodesic flow on modular surfaces and continued fractions*, Ann. Sci. École Norm. Sup. (4) **26** (1993), no. 1, 23–50.

[KM96] Vadim A. Kaimanovich and Howard A. Masur, *The Poisson boundary of the mapping class group*, Invent. Math. **125** (1996), no. 2, 221–264.

[Ker80] Steven P. Kerckhoff, *The asymptotic geometry of Teichmüller space*, Topology **19** (1980), no. 1, 23–41.

[Kes59] Harry Kesten, *Symmetric random walks on groups*, Trans. Amer. Math. Soc. **92** (1959), 336–354.

[Khi97] A. Ya. Khinchin, *Continued fractions*, Russian, Dover Publications Inc., Mineola, NY, 1997. With a preface by B. V. Gnedenko, Reprint of the 1964 translation.

[Mah10a] Joseph Maher, *Linear progress in the complex of curves*, Trans. Amer. Math. Soc. **362** (2010), no. 6, 2963–2991.

[Mah10b] Joseph Maher, *Random Heegaard splittings*, J. Topol. **3** (2010), no. 4, 997–1025.

[Mah11a] Joseph Maher, *Random walks on the mapping class group*, Duke Math. J. **156** (2011), no. 3, 429–468.

[Mah11b] Joseph Maher, *Exponential decay in the mapping class group* (2011), available at `arXiv:1104.5543`.

[Mas85] Bernard Maskit, *Comparison of hyperbolic and extremal lengths*, Ann. Acad. Sci. Fenn. Ser. A I Math. **10** (1985), 381–386.

[Mas80] Howard A. Masur, *Uniquely ergodic quadratic differentials*, Comment. Math. Helv. **55** (1980), no. 2, 255–266.

[Mas82] Howard A. Masur, *Interval exchange transformations and measured foliations*, Ann. of Math. (2) **115** (1982), no. 1, 169–200.

[Mas93] Howard A. Masur, *Logarithmic law for geodesics in moduli space*, (Göttingen, 1991/Seattle, WA, 1991), Contemp. Math., vol. 150, Amer. Math. Soc., Providence, RI, 1993, pp. 229–245.

[MM99] Howard A. Masur and Yair N. Minsky, *Geometry of the complex of curves. I. Hyperbolicity*, Invent. Math. **138** (1999), no. 1, 103–149.

[MM00] Howard A. Masur and Yair N. Minsky, *Geometry of the complex of curves. II. Hierarchical structure*, Geom. Funct. Anal. **10** (2000), no. 4, 902–974.

[MS91] Howard A. Masur and John Smillie, *Hausdorff dimension of sets of nonergodic measured foliations*, Ann. of Math. (2) **134** (1991), no. 3, 455–543.

[Min96a] Yair N. Minsky, *Extremal length estimates and product regions in Teichmüller space*, Duke Math. J. **83** (1996), no. 2, 249–286.

[Min96b] Yair N. Minsky, *Quasi-projections in Teichmüller space*, J. Reine Angew. Math. **473** (1996), 121–136.

[Raf07] Kasra Rafi, *A combinatorial model for the Teichmüller metric*, Geom. Funct. Anal. **17** (2007), no. 3, 936–959.

[Raf10] Kasra Rafi, *Hyperbolicity in Teichmüller space* (2010), available at `arXiv:1011.6004`.

[Ser81] Caroline Series, *On coding geodesics with continued fractions*, Ergodic theory (Sem., Les Plans-sur-Bex, 1980), Monograph. Enseign. Math., vol. 29, Univ. Genève, Geneva, 1981, pp. 67–76.

[Str84] Kurt Strebel, *Quadratic differentials*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 5, Springer-Verlag, Berlin, 1984.

[Sul82] Dennis Sullivan, *Disjoint spheres, approximation by imaginary quadratic numbers, and the logarithm law for geodesics*, Acta Math. **3-4** (1982), 215–237.

[Tio12] Giulio Tiozzo, *Sublinear deviation between geodesics and sample paths* (2012), available at `arXiv:1210.7352`.

[Vor05] Yaroslav Vorobets, *Periodic geodesics on generic translation surfaces*, Contemp. Math., vol. 385, Amer. Math. Soc., Providence, RI, 2005.

Vaibhav Gadre
Harvard University
`vaibhav@math.harvard.edu`

Joseph Maher
CUNY College of Staten Island and CUNY Graduate Center
`joseph.maher@csi.cuny.edu`

Giulio Tiozzo
Harvard University
`tiozzo@math.harvard.edu`